

CONTRIBUTION TO TABANIDAE FAUNA OF WEST AEGEAN REGION (INSECTA: DIPTERA)

Ferhat Altunsoy* and A. Yavuz Kılıç*

* Anadolu University, Faculty of Science, Department of Biology, 26470 Eskişehir / TURKEY.

[Altunsoy, F. & Kılıç, A. Y. 2016. Contribution to Tabanidae fauna of West Aegean Region (Insecta: Diptera). Munis Entomology & Zoology, 11 (1): 223-229]

ABSTRACT: This study is conducted to determine West Aegean Region Tabanidae fauna in 2012 to 2013. As a result 3 subfamilies, 10 genera and 52 species are determined. 16 species are firstly reported from region. These species are *Atylotus loewianus* Villeneuve, 1920, *Therioplectes tunicatus* Szilady, 1927, *Hybomitra caucasi* Szilady, 1923, *Tabanus cordiger* Meigen, 1820, *Tabanus fraseri* Austen, 1924, *Tabanus maculicornis* Zetterstedt, 1842, *Tabanus miki* (Brauer, 1880), *Tabanus nemoralis* Meigen, 1820, *Tabanus oppugnator* Austen, 1925, *Tabanus portschinskii* Olsufjev, 1937, *Tabanus spodopteroideus* Olsufjev, Moucha & Chvála, 1969, *Tabanus sudeticus* Zeller 1842, *Tabanus tergestinus* Egger, 1859, *Tabanus unifasciatus* Loew, 1858, *Haematopota italica* Meigen, 1804, *Philipomyia aprica* (Meigen, 1820). As a conclusions number of species which are distributing in the region reach to 67.

KEY WORDS: West Aegean, Fauna, Horse fly, Tabanidae, Diptera, Turkey

Female horse fly species during the blood-feeding period frequently change hosts in mammals including human. They have great importance in terms of medical and veterinary, because they are potential mechanical vectors for the diseases caused by viruses, bacteria and protozoans and the economic significance of stress resulting directly from bites, or indirect secondary infections such as anemia through blood loss, allergic responses, etc. (Chvala et al., 1972; Krinsky, 1979; Foil, 1989).

Studies about Tabanidae fauna of Turkey have begun in the 19th century (Walker, 1854; Loew, 1856a,b,c, 1858, 1859) and still continue. Many studies were reported about distribution, seasonality and habitat preferences of Turkish horse fly species (Kılıç, 1992, 1993, 1996a,b,c, 1999, 2001a,b,c, 2002, 2003, 2004, 2005a,b, 2006; Kılıç & Schacht, 1995; Altunsoy & Kılıç, 2011a,b, 2012, 2014; Altunsoy, 2011). Nevertheless these studies are not adequate for the put forth of Turkish horse fly fauna.

Based on reports of recent studies, Turkish horse fly fauna is representing with 3 subfamilies, 9 genera, 171 species and 15 subspecies (Kılıç, 1999, 2006; Altunsoy & Kılıç, 2014) and previously 51 species were reported from Aegean region (Schacht, 1984, 1985, 1987). In this study, totally 1672 samples were collected and 52 species were identified and 16 species were reported for the first time from the study area. As a result, the number of species which are distributed in Aegean region was reached 67.

MATERIAL AND METHODS

Adult female horse flies were collected from different habitats in West Aegean Region (Aydın, Denizli, İzmir, Manisa and Muğla) with Malaise and Nzi Traps, which were baited with 1-octen-3-ol, and water traps.

Collection and preparation of samples were done according to the principles of Chvala et al. (1972) and Olsufjev (1977). Tabanids were killed by ethyl-acetat jars. The specimens were brought to the laboratory in 70 degree alcohol solution and were pinned with insect pins.

Samples were identified according to Chvala et al. (1972), Olsufjev (1977), Peus (1980), Yücel (1987), Schacht (1987), Leclercq (1966a,b, 1967a,b) and Rubio (2002). Identified samples were preserved in the Zoological Museum of Anadolu University (AUZM).

The distributions of species in Turkey and worldwide because of given in previous studies, not presented here again (Kılıç, 1999, 2006; Schacht, 1983, 1984, 1985, 1987; Andreeva et al., 2009; Altunsoy & Kılıç, 2010).

RESULTS AND DISCUSSION

Totally 52 species belonging to 9 genera and 3 subfamilies were identified and 16 species were recorded for the first time from Aegean Region.

Family TABANIDAE

Subfamily PANGONINAE

Tribe Pangoniini

Pangonius fulvipes (Loew, 1859)

Material examined: İzmir (Kemalpaşa), 10.05.2012, 2♀♀.

Pangonius pyritosus Loew, 1859

Material examined: İzmir (Kemalpaşa), 10.05.2012, 2♀♀.

Subfamily CHRYSOPSINAE

Tribe Chrysopsini

Chrysops caecutiens (Linne, 1761)

Material examined: Muğla (Fethiye-Köyceğiz), 07.05.2012, 5♀♀, 1♂; Muğla (Marmaris), 10.06.2012, 1♀; Uşak (Karaağaç), 15.06.2012, 10♀♀; Uşak (Central), 12.06.2012, 3♀♀; Uşak (Çamyuva), 15.06.2012, 4♀♀; İzmir (Kiraz), 13.07.2012, 13♀♀.

Chrysops flavipes (Linne, 1761)

Material examined: Uşak (Güre), 09.06.2013, 3♀♀; 11.06.2013, 12♀♀; Uşak (Derbent), 09.09.2013, 2♀♀; Marmaris (Çamlı), 16.05.2013, 3♀♀.

Silvius alpinus Scopoli, 1763

Material examined: Manisa (Demirci), 07.07.2013, 1♀.

Subfamily TABANINAE

Tribe Tabanini

Atylotus fulvus (Meigen, 1820)

Material examined: Denizli (Buldan), 14.07.2012, 2♀♀.

Atylotus loewianus Villeneuve, 1920

Material examined: Denizli (Buldan), 14.07.2012, 1♀.

Therioplectes tricolor Zeller, 1842

Material examined: Muğla (Fethiye-Köyceğiz), 07.05.2012 2♀♀; Muğla (Marmaris), 07.05.2012, 1♀; 10.06.2013, 2♀♀; Marmaris (Çamlı), 16.05.2013, 2♀♀; Kuşadası, 12.05.2013 2♀♀; Uşak (Güre), 16.05.2013, 1♀.

Therioplectes tunicatus Szilady, 1927

Material examined: Muğla (Dalaman), 16.05.2013, 1♀.

Hybomitra acuminata (Loew, 1858)

Material examined: Denizli (Central), 25.05.2012, 5♀♀, Denizli (Buldan), 15.05.2013, 8♀♀.

Hybomitra caucasi Szilady, 1923

Material examined: Denizli (Central), 25.05.2012, 4♀♀, 2♂♂.

Hybomitra caucasica (Enderlein, 1925)

Material examined: Manisa (Central), 23.05.2012, 3♀♀.

***Hybomitra ciureai* (Séguy, 1937)**

Material examined: Denizli (Buldan), 14.07.2012, 15♀, 2 ♂; Muğla (Fethiye), 16.05.2013, 2♀; Muğla (Dalaman), 16.05.2013, 3♀; Muğla (Köyceğiz), 16.05.2013, 5♀; Muğla (Marmaris), 10.06.2012, 1♀.

***Tabanus autumnalis* (Linne, 1761)**

Material examined: Muğla (Fethiye-Köyceğiz), 07.05.2012, 1♀ 11.05.2013, 1♀, 16.05.2013, 1♀; Muğla (Köyceğiz), 16.05.2013, 1♀; Muğla (Datça), 11.06.2012, 1♀; Muğla (Dalaman), 16.05.2013, 1♀; Uşak (Karaağaç), 15.06.2012, 4♀, 11.05.2013 2♀ ; Uşak (Ulubey), 09.07.2012, 2♀; Uşak (Eşme), 17.05.2013, 1♀; Uşak (Central), 20.07.2012, 1♀; Denizli (Çal), 10.07.2012, 3♀; Denizli (Buldan), 14.07.2012, 2♀; 14.05.2013, 8 ♀.

***Tabanus bifarius* Loew, 1858**

Material examined: Denizli (Cankurtaran), 09.06.2012, 15♀ 12.05.2013 5♀, 12.08.2013 3♀; Denizli (Pamukkale), 08.06.2012, 7♀; Uşak (Central), 11.06.2012, 4♀; 20.07.2012, 6♀; Uşak (Eşme), 09.06.2013, 2♀; 20.07.2012 1♀; 17.05.2013 1♀; Uşak (Çamyuva), 15.06.2012, 4♀; Uşak (Güre), 09.06.2013, 4♀; Uşak (Karaağaç), 15.06.2012, 1♀; Manisa (Central), 11.06.2012, 3♀ 10.06.2013 3♀; Manisa (Demirci), 07.07.2013, 1♀; Denizli (Honaz Dağı), 12.06.2012, 5♀; 10.06.2013, 4♀; Muğla (Fethiye-Köyceğiz), 07.05.2012, 4♀; 05.06.2012 2♀; 16.05.2013 1♀; Muğla (Marmaris), 07.05.2012, 1♀; 10.06.2012, 9♀; 11.05.2013 2♀, 13.08.2013 2♀, 11.06.2013, 4♀; Marmaris (Değirmen), 16.05.2013, 1♀; Muğla (Fethiye), 16.05.2013, 4♀; Muğla (Köyceğiz), 16.05.2013, 1♀; Muğla (Datça), 11.06.2012, 2♀; Muğla (Değirmen), 16.05.2013, 1♀; Kuşadası, 15.05.2013, 14♀; Aydın (Karacasu), 16.05.2013, 1♀; İzmir (Ödemiş, Bozdağ), 08.07.2013, 2♀.

***Tabanus bromius* Linne, 1761**

Material examined: Uşak (Central), 14.07.2012, 3♀; 10.06.2013, 4♀; 07.07.2013, 4♀; 13.08.2013, 6♀; Muğla (Fethiye-Köyceğiz), 11.08.2012, 8♀, 08.07.2013 5♀, 14.08.2013, 6♀; İzmir (Ödemiş), 12.08.2012, 5♀, 10.06.2013, 4♀, Manisa (Salihli), 12.08.2012, 23♀; Denizli (Buldan), 14.07.2012, 17♀; İzmir (Kiraz), 13.07.2012, 2 3♀; Denizli (Honaz Dağı), 10.07.2012, 2♀ 12.06.2013 5♀; Denizli (Honaz), 10.07.2012, 1♀; Denizli (Çameli), 10.07.2012, 8♀; Muğla (Ula), 11.07.2012, 18♀, 06.07.2013 10♀; İzmir (Beydağı), 13.07.2012, 2♀; Uşak (Central), 11.06.2012, 4♀; 20.07.2012, 1♀; Uşak (Karaağaç), 15.06.2012, 10♀, 16.08.2013 7♀; Uşak (Çamyuva), 15.06.2012, 14 ♀; 16.06.2013, 10♀, 11.05.2013, 9♀; Muğla (Milas), 10.06.2012, 5♀, 11.06.2013, 2♀; Denizli (Honaz Dağı), 12.06.2012, 10♀; Uşak (Ulubey), 09.07.2012, 5♀; Denizli (Çal), 10.07.2012, 5♀, Kuşadası 12.05.2013, 2♀; Aydın (Horsunlu), 10.06.2013, 8♀; Muğla (Ören), 12.06.2013, 8 ♀; 09.07.2013, 12 ♀; Muğla (Marmaris), 10.06.2012, 1♀; Muğla (Kale), 12.09.2013, 1♀; Muğla (Fethiye, Çameli Yolu), 10.05.2012, 3♀.

***Tabanus cordiger* Meigen, 1820**

Material examined: Uşak (Karaağaç), 15.06.2012, 3♀; Manisa (Central), 11.06.2012, 3♀; İzmir (Spil Dağı), 13.07.2013, 2♀; Muğla (Marmaris), 10.06.2012, 3♀; Muğla (Fethiye, Çameli Yolu), 10.05.2012, 1♀.

***Tabanus exclusus* Pandelle, 1883**

Material examined: Denizli (Buldan), 14.07.2012, 1♀; Denizli (Honaz), 10.07.2012, 13♀; Denizli (Buldan), 14.07.2012, 1♀; Uşak (Central), 20.07.2012, 1♀; Muğla (Fethiye, Çameli Yolu), 10.05.2012, 21♀; Muğla (Marmaris-Fethiye, Ula Yolu), 11.07.2012, 6♀; Muğla (Fethiye-Köyceğiz), 25.04.2012, 3♀.

***Tabanus fraseri* Austen, 1924**

Material examined: Uşak (Derbent), 09.09.2013, 8♀; Muğla (Kavaklıdere), 11.09.2013, 1♀; İzmir (Ödemiş, Bozdağ), 01.09.2013, 2♀; İzmir (Beydağ), 11.09.2013, 1♀; Aydın (Bazdoğan), 11.09.2013, 1♀; Muğla (Central), 12.09.2013, 2♀; İzmir (Beydağ, Çamedli-Alısu), 11.09.2013, 1♀.

***Tabanus glaucopis* Meigen, 1936**

Material examined: İzmir (Dikili), 16.07.2012, 5♀; Denizli (Buldan) 10.07.2012 15♀; 14.07.2012, 4♀; Uşak (Central), 12.07.2012, 11♀; 20.07.2012, 2♀; Aydın (Nazilli), 12.07.2013, 11♀.

***Tabanus indrae* Hauser, 1939**

Material examined: Muğla (Datça), 17.06.2012, 4♀.

***Tabanus leleani* Austen, 1920**

Material examined: Uşak (Derbent), 09.09.2013, 2♀.

***Tabanus laetitinctus* Becker, 1912**

Material examined: Muğla (Datça), 17.06.2012, 6 ♀♀.

***Tabanus lunatus* Fabricius, 1794**

Material examined: Muğla (Ortaca), 11.07.2012, 15♀, 13.07.2013 10♀; Deniz (Kale), 12.07.2012, 6♀, Uşak (Central), 14.07.2012, 5♀; 20.07.2012, 2♀; İzmir (Kiraz), 13.07.2012, 32♀, Denizli (Honaz Dağı), 10.07.2012, 25♀; Denizli (Honaz), 10.07.2012, 3♀; Denizli (Çameli), 10.07.2012, 30♀; Muğla (Ula), 11.07.2012, 14♀; Aydın (Karacasu), 13.07.2012, 10♀, İzmir (Beydağı), 13.07.2012, 5♀, Muğla (Milas), 10.06.2012, 5♀, 10.06.2013 6♀, 09.07.2013 4♀; 11.06.2013, 3♀; Muğla (Marmaris), 10.06.2013, 7♀; Muğla (Fethiye, Çameli Yolu), 10.05.2012, 16♀; Denizli (Çal), 10.07.2012, 18♀; Muğla (Marmaris-Fethiye, Ula Yolu), 11.07.2012, 5♀; Manisa (Demirci), 07.07.2013, 3♀.

***Tabanus maculicornis* Zettersted, 1842**

Material examined: Muğla (Marmaris, Değirmen), 16.05.2013, 1♀.

***Tabanus miki* (Brauer, 1880)**

Material examined: Denizli (Kiraz), 13.07.2012, 1♀; Denizli (Honaz Dağı), 10.07.2012, 8♀; Denizli (Honaz), 10.07.2012, 4♀; Denizli (Buldan), 14.07.2012, 2♀; Aydın (Horsunlu), 10.06.2013, 2♀; İzmir (Beydağı), 13.07.2012, 5♀; İzmir (Beydağ, Çamedli-Alısu), 11.09.2013, 1♀; Uşak (Central), 14.07.2012, 3♀; 20.07.2012, 1♀; Muğla (Fethiye-Köyceğiz), 11.08.2012, 8♀; İzmir (Ödemiş), 12.08.2012, 15♀, Manisa (Salihli), 12.08.2012, 5♀; Muğla (Fethiye, Çameli Yolu), 10.05.2012, 1♀.

***Tabanus nemoralis* Meigen, 1820**

Material examined: Uşak (Eşme), 09.06.2013, 1♀.

***Tabanus obsolescens* (Pandelle, 1883)**

Material examined: Manisa (Turgutlu) 15.06.2012, 4♀; Denizli (Buldan) 12.06.2012 2♀; Denizli (Honaz), 10.07.2012, 2♀; İzmir (Ödemiş, Bozdağ), 01.09.2013, 4♀; 11.09.2013, 1♀; İzmir (Beydağ), 11.09.2013, 1♀; İzmir (Beydağ, Çamedli-Alısu), 11.09.2013, 17♀; Muğla (Central), 12.09.2013, 43♀; Muğla (Kale), 12.09.2013, 21♀; Muğla (Kavaklıdere), 11.09.2013, 27♀; Aydın (Bazdoğan), 11.09.2013, 21♀; Uşak (Derbent), 09.09.2013, 3♀; Muğla (Milas, Labranda), 09.07.2013, 1♀.

***Tabanus oppugnator* Austen, 1925**

Material examined: Uşak (Güre), 08.06.2013, 2♀.

***Tabanus prometheus* Szilady, 1923**

Material examined: Uşak (Central), 11.06.2012, 2♀.

***Tabanus portschinskii* Olsufjev, 1937**

Material examined: Uşak (Central), 15.07.2012, 8♀; Denizli (Buldan) 10.07.2012 2♀.

***Tabanus quatuornotatus* Meigen, 1820**

Material examined: Denizli (Pamukkale), 06.05.2012, 8♀, 1♂; 08.06.2012, 4♀; Denizli (Honaz), 10.05.2012, 5♀; Muğla (Datça-Kinidos), 07.05.2012, 6♀; Muğla (Marmaris-Knidos), 09.05.2012, 5♀; Muğla (Dalaman), 16.05.2013, 1♀; Muğla (Datça), 11.06.2012, 2♀; Muğla (Marmaris), 10.06.2012, 5♀; Denizli (Central), 25.05.2012, 5♀, 2♂♂; İzmir (Spil Dağı), 14.06.2012, 9♀; Aydın (Kuşadası), 12.05.2013, 5♀; Aydın (Karacasu), 16.05.2013, 12♀; Uşak (Eşme), 17.05.2013, 7♀; 20.07.2012, 6♀; Uşak (Central), 11.06.2012, 3♀; 20.07.2012, 12♀; Uşak (Güre), 09.06.2013, 1♀; Uşak (Karaağaç), 15.06.2012, 1♀.

***Tabanus regularis* Jaennicke, 1866**

Material examined: İzmir (Kemalpaşa), 10.05.2012, 2♀; Muğla (Fethiye-Köyceğiz), 07.05.2012, 2♀; Denizli (Kiraz), 13.07.2012, 1♀; Muğla (Milas, Labranda), 09.07.2013, 4♀; Muğla (Fethiye, Çameli Yolu), 10.05.2012, 3♀; Muğla (Gökova), 11.07.2013, 1♀.

***Tabanus rupium* (Brauer & Bergenstamm, 1880)**

Material examined: Denizli (Honaz Dağı), 10.07.2012, 1♀; Denizli (Honaz), 10.07.2012, 1♀; Uşak (Karaağaç), 15.06.2012, 5♀; Uşak (Çamyuva), 15.06.2012, 4♀; Uşak (Eşme), 17.05.2012, 2♀; 20.07.2012, 16♀; Muğla (Marmaris), 10.06.2012, 2♀; Muğla (Datça), 11.06.2012, 2♀; Uşak (Central), 20.07.2012, 6♀; Aydın (Karacasu), 16.05.2013, 1♀; Uşak (Güre), 09.06.2013, 1♀; Uşak (Güney), 17.05.2013, 1♀.

***Tabanus spodopterus* Meigen, 1820**

Material examined: Denizli (Honaz Dağı), 10.07.2012, 4♀; Denizli (Çameli), 10.07.2012, 4♀; Denizli (Buldan) 12.07.2012 12♀. Muğla (Ula), 11.07.2012, 4♀; Muğla (Ortaca), 11.07.2012, 2♀; Denizli (Kale), 12.07.2012, 1♀; Aydın (Karacasu), 13.07.2012, 4♂; İzmir (Beydağı), 13.07.2012, 2♀; İzmir (Kiraz), 13.07.2012, 6♀; Uşak (Central), 14.07.2012, 2♀; Muğla (Fethiye-Köyceğiz), 11.08.2012, 2♀; Muğla (Marmaris-Fethiye, Ula), 2♀; Muğla (Fethiye-Köyceğiz), 20.04.2012, 1♀; Muğla (Fethiye, Çamedli), 2♀; Muğla (Milas, Labranda), 09.07.2013, 1♀.

***Tabanus spodopteroides* Olsufjev, Moucha & Chvála, 1969**

Material examined: Denizli (Honaz Dağı), 10.07.2012, 11♀; Denizli (Çameli), 10.07.2012, 4♀; Uşak (Güre), 09.06.2013, 1♀; Muğla (Fethiye-Çameli Yolu), 10.05.2012, 1♀.

***Tabanus sudeticus* Zeller 1842**

Material examined: Denizli (Buldan) 12.07.2012 2♀.

***Tabanus tunicatus* Szilady, 1927**

Material examined: Muğla (Fethiye) 17.05.2013, 2♀.

***Tabanus tinctus* (Walker, 1850)**

Material examined: Denizli (Buldan) 12.07.2012 2♀; Aydın (Karacasu), 13.07.2012, 4♂; 2♀; Uşak (Central), 14.07.2012, 3♀.

***Tabanus tergestinus* Egger, 1859**

Material examined: Muğla (Köyceğiz), 07.05.2012, 15♀; 16.05.2013, 1♀; Muğla (Fethiye) 15.05.2013, 10♀; Kuşadası, 15.05.2013, 14♀.

***Tabanus unifasciatus* Loew, 1858**

Material examined: Uşak (Çamyuva), 15.06.2012, 4♀; Uşak (Karaağaç), 15.06.2012, 6♀; Uşak (Central), 20.07.2012, 7♀; Muğla (Fethiye) 15.05.2013, 4♀; Muğla (Datça), 11.06.2012, 2♀; Uşak (Eşme), 20.07.2012, 2♀; Muğla (Marmaris), 10.06.2012, 4♀.

Tribe Haematopotini***Haematopota bigoti* Gobert, 1880**

Material examined: İzmir (Dikili), 16.07.2012 1♀.

***Haematopota long antennata* (Olsufjev 1937)**

Material examined: Aydın (Karacasu), 10.06.2013, 2♀.

***Haematopota grandis* Meigen 1820**

Material examined: Muğla (Datça), 17.06.2012, 6♀.

***Haematopota ocelligera* (Krober 1922)**

Material examined: Muğla (Fethiye), 08.06.2013 9♀.

***Haematopota pallens* Loew 1871**

Material examined: İzmir (Dikili), 16.07.2012 2♀.

***Haematopota subcylindrica* Pandelle, 1883**

Material examined: Uşak (Central), 12.06.2012, 5♀.

***Haematopota italica* Meigen, 1804**

Material examined: Muğla (Fethiye- Köyceğiz), 07.05.2012, 5♀, Muğla (Marmaris), 07.05.2012, 1♀.

Tribe Diachlorini

Dasyrhamphis carbonarius (Meigen, 1820)

Material examined: Muğla (Marmaris), 07.05.2012, 1♀; 10.06.2012, 2♀♀; Marmaris (Çamlı), 16.05.2013, 2♀♀; Muğla (Dalaman), 16.05.2013, 1♀; Denizli (Honaz Dağı), 12.06.2012, 2♀♀; Uşak (Central), 12.06.2012, 2♀♀; 14.06.2013, 4♀♀; Uşak (Eşme), 20.07.2012, 1♀; Aydın (Karacasu), 10.06.2013, 2♀♀.

Dasyrhamphis umbrinus (Meigen, 1820)

Material examined: Denizli (Pamukkale), 06.05.2012, 1♀, Muğla (Fethiye- Köyceğiz), 07.05.2012, 2♀♀; Muğla (Marmaris), 07.05.2012, 6♀♀; Muğla (Köyceğiz), 16.05.2013, 2♀♀; Muğla (Fethiye), 16.05.2013, 2♀♀; Marmaris (Çamlı), 16.05.2013, 1♀; Muğla (Datça-Kinidos), 07.05.2012, 1♀; Muğla (Dalaman), 16.05.2013, 1♀; Denizli (Honaz Dağı), 12.06.2012, 5♀♀; İzmir (Spil Dağı), 14.06.2012, 7♀♀; Aydın (Kuşadası), 12.05.2013, 1♀; Uşak (Güney), 17.05.2013, 1♀.

Philipomyia aprica (Meigen, 1820)

Material examined: Uşak (Central), 15.07.2012, 12♀♀; Uşak (Güre), 09.06.2013, 6♀♀; Kuşadası, 15.05.2013, 12♀♀; Muğla (Fethiye), 16.05.2013, 1♀; 08.06.2013 9♀♀, 6♂♂; İzmir (Ödemiş, Bozdağ), 08.07.2013, 9♀♀.

Philipomyia graeca (Fabricius, 1794)

Material examined: Muğla (Fethiye- Köyceğiz), 07.05.2012, 12♀♀; 13.06.2013, 22♀♀; Muğla (Marmaris), 07.05.2012, 12♀♀; Marmaris (Çamlı), 16.05.2013 2♀♀; Muğla (Datça-Kinidos), 07.05.2012, 3♀♀, Muğla (Milas), 10.06.2012, 25♀♀, Milas (Labranda), 11.06.2013, 2♀♀; Muğla (Köyceğiz), 16.05.2013, 5♀♀; Muğla (Dalaman), 16.05.2013, 2♀♀.

In the previous studies, a total of 51 species were reported from Aegean Region. 16 species are firstly reported from this region with this work: *Atylotus loewianus* Villeneuve, 1920, *Hybomitra caucasi* Szilady, 1923, *Tabanus cordiger* Meigen, 1820, *Tabanus fraseri* Austen, 1924, *Tabanus miki* (Brauer, 1880), *Tabanus maculicornis* Zetterstedt, 1842, *Tabanus nemoralis* Meigen, 1820, *Tabanus oppugnator* Austen, 1925, *Tabanus portschinskii* Olsufjev, 1937, *Tabanus spodopteroideus* Olsufjev, Moucha & Chvála, 1969, *Tabanus sudeticus* Zeller 1842, *Tabanus tergestinus* Egger, 1859, *Tabanus tunicatus* Szilady, 1927 *Tabanus unifasciatus* Loew, 1858 ve *Haematopota italica* Meigen, 1804.

Results of study indicated that the *Tabanus bromius* as the most abundant species with 18%. It was determined in previous studies that *Tabanus bromius* most abundant species in the any habitats (Yücel, 1987; Kılıç, 1992, 2001c, 2004, 2005b). This species is followed by *Tabanus lunatus* (13.3%) and *Tabanus obsolescens* (8.8%). These three species consist of 40.1% of the horse fly fauna on the study area. This study does not contain all species for West Aegean Region, but shows the importance of periodic studies and isolated areas for faunistic studies. In addition, it can be inferred that the species which known as unique can be observed in many different areas.

LITERATURE CITED

- Altunsoy, F. & Kılıç, A. Y. 2010. A New Record for Turkish Tabanidae (Insecta: Diptera) Fauna. J. Ent. Res. Soc., 12(2): 109-111.
- Altunsoy, F. & Kılıç, A. Y. 2011a. New Data About *Tabanus karaosus* Timmer 1984 (Diptera: Tabanidae) from Turkey. J. Ent. Res. Soc., 13 (1): 75-80.
- Altunsoy, F. & Kılıç, A. Y. 2011b. Doğu Karadeniz Bölgesi Tabanidae (Diptera: Insecta) Faunası. Gümmüşhane Üniversitesi, Fen Bilimleri Enstitüsü Dergisi, 1 (1): 24-36.
- Altunsoy, F. & Kılıç, A. Y. 2012. Seasonal Abundance of Horse Fly (Diptera: Tabanidae) in Western Anatolia. J. Entomol. Res. Soc., 14 (1): 95-105.
- Altunsoy, F. & Kılıç, A. Y. 2014. New records for Turkish Tabanidae (Insecta: Diptera) Fauna. Turkish Journal of Entomology, (In press).
- Andreeva, R., Altunsoy, F. & Kılıç, A. Y. 2009. New Contribution to Information about Tabanidae (Diptera) Adult and Larvae from West Anatolia. J. Ent. Res. Soc., 11 (3): 19-30.
- Chvala, M., Lyneborg, L. & Moucha, J. 1972. The horse flies of Europe (Diptera: Tabanidae). Ent. Soc. Copenhagen, E. W. Classey Ltd. Hampton, 453 pp.
- Foil, L. 1989. Tabanids as vectors of disease agents. Parasitology Today, 5: 88-96.

- Kılıç, A. Y.** 1992. Eskişehir ve Çevresi Tabanidae (Diptera) Faunasının İncelenmesi. *Türk Entomoloji Dergisi*, 16 (3): 169-180.
- Kılıç, A. Y.** 1993. Eskişehir ve Çevresi *Tabanus bromius* L., *T. exlusus* Pand., *T. glaucopsis* Meig., *T. spodopterus ponticus* Ols., Mch. Chv., ve *Philipomyia aprica* (Meig) (Diptera: Tabanidae) türlerinin mevsimsel aktiviteleri üzerine bir çalışma. *Doğa Türk Zooloji Dergisi*, 17 (3): 303-310.
- Kılıç, A. Y.** 1996a. Çamlıyayla (İçel) Tabanidae (Diptera) faunası üzerine araştırmalar. *Türk Entomoloji dergisi*, 20 (2): 123-135.
- Kılıç, A. Y.** 1996b. Bilecik ili Tabanidae (Diptera) Faunası üzerine araştırmalar. *Anadolu Üni. Fen Fak. Derg.*, 1: 45-57.
- Kılıç, A. Y.** 1996c. Türkiye Tabanidae (Diptera) Faunası için iki yeni kayıt ve bazı türlerin yeni lokalite kayıtları, *Anadolu Üni. Fen Fak. Derg.*, 2: 105-115.
- Kılıç, A. Y.** 1999a. Checklist of Tabanidae (Diptera) From Turkey. *Tr. J. Zool.*, 23 (2): 123-132.
- Kılıç, A. Y.** 1999b. Trakya Bölgesi Tabanidae (Diptera) Faunası. *Türk Zooloji dergisi*, 23 (1): 67-89.
- Kılıç, A. Y.** 2001a. The Tabanidae (Diptera) Fauna of Balıkesir Province. *Türk J. Zool.*, 25: 395-402.
- Kılıç, A. Y.** 2001b. The Tabanidae (Diptera) Fauna of Çanakkale Province. *Türk J. Zool.*, 25: 403-411.
- Kılıç, A. Y.** 2001c. The Tabanidae (Diptera) Fauna of Kütahya Province of Turkey. *J. Ent. Res. Soc.*, 3 (3): 29-41.
- Kılıç, A. Y.** 2003. Bursa ve Yalova İlleri Tabanidae (diptera) faunası Üzerinde Araştırmalar. *Türk Ent. Derg.*, 27 (3): 207-221.
- Kılıç, A. Y.** 2004. Bolu ili Tabanidae (Insecta: Diptera) Faunası. *Türkiye Entomoloji dergisi*, 28 (1): 57-68.
- Kılıç, A. Y.** 2005a. Sinop ili Tabanidae (Insecta: Tabanidae) Faunası. *Anadolu Üni. Bilim ve Teknoloji Derg.*, 5 (2): 207-221.
- Kılıç, A. Y.** 2005b. Bartın, Karabük ve Zonguldak illeri Tabanidae (Diptera) faunası üzerinde araştırmalar. *Türk Ent. Derg.*, 29 (2): 51-60.
- Kılıç, A. Y.** 2006. New addition and errata to the checklist of Tabanidae (Insecta: Diptera) fauna of Turkey. *Turk. J. Zool.*, 30 (2): 335-343.
- Kılıç, A. Y. & Öztürk, R.** 2002. Sultandağı çevresinin Tabanidae (Diptera) Faunası üzerine çalışmalar. *Anadolu Üni. Bilim ve Teknoloji Dergisi*, 3 (2): 307-316.
- Kılıç, A. Y. & Schacht, W.** 1995. Eine neue Bremsenart aus der West-Türkei (Diptera: Tabanidae). *Entomofauna*, 16 (10): 245-252.
- Kılıç, A. Y., Gören, T. & Altunsoy, F.** 2010. Düzce ili Tabanidae (Insecta: Diptera) faunası. *Sakarya Üniversitesi, Fen Bilimleri Dergisi*, 14 (1): 29-32.
- Krinsky, W. L.** 1979. Animal disease agents transmitted by horse flies and deer flies (Diptera: Tabanidae). *Journal of Medical Entomology*, 13: 225-275.
- Leclercq, M.** 1966a. Revision systematique et biogeographique des Tabanidae (Diptera) Palearctiques, Tabaninae. *Mem. Ins. Roy. Sci. Nat. Belg.*, II (79): 1-236.
- Leclercq, M.** 1966b. Tabanidae (Diptera) de Turquie diagnosis d'*Atylotus hendrixi*, *Haematopota coolsi*, *Haematopota delozi* n. spp. *Bull. Rech. Agron. Gembloux*, 1 (3): 463-477.
- Leclercq, M.** 1967a. Tabanidae (Diptera) de Turquie, II. Diagnosis D'*Hybomitra okayi*, *Atylotus hendrixi*, et *Haematopota hennauxi* n. spp. *Bull. Rech. Agron. Gembloux*, 2 (1): 106-127.
- Leclercq, M.** 1967b. Tabanidae (Diptera) de Turquie III. *Bull. Rech. Agron. Gembloux*, 2 (4): 707-710.
- Loew, H.** 1856a. Neue Beiträge zur Kenntnis der Dipteren, 4. Beitrag, Berlin, 57 pp. Tabanidae 24-28 pp.
- Loew, H.** 1856b. Zur Kenntnis der Europäischen *Tabanus* Arten. *Verh. Zool. Bot. Ges.W. en.*, 8: 573-612.
- Loew, H.** 1856c. Versuch einer Auseinander setzung der Europäischen Chrysops Arten. *Ibid.*, 8: 613-634.
- Loew, H.** 1858. Ueber die Europäischen Arten der Gattung *Silvius*. *Ibid.*, 2: 350-352.
- Loew, H.** 1859. Neue Beiträge zur Kenntnis der Dipteren, 6. Beitrag, Berlin, 50 pp. Tabanidae 23-32 pp.
- Olsufjev, N. G.** 1977. Faune de l'URSS insectes Dipteres, VII, 2: Tabanidae. *Acad. Sci. URSS. Trav. Zool.* 113, Leningrad, 434 pp.
- Schacht, W.** 1983. Eine neue Bremsenart aus der Türkei (Diptera: Tabanidae). *Entomofauna*, 4 (27): 483-492.
- Schacht, W.** 1984. Beitrag zu einigen paläarktischen bremsen Arten vornehmlich aus der Türkei (Diptera: Tabanidae). *Entomofauna*, 5 (35): 483-498.
- Schacht, W.** 1985. Kleiner Beitrag zur bremsen fauna der Türkei (Diptera: Tabanidae). *Entomofauna*, 6 (28): 501-508.
- Schacht, W.** 1987. Ein weiterer Beitrag bremsen fauna der Türkei (Diptera: Tabanidae). *Entomofauna*, 8 (33): 485-496.
- Walker, F.** 1854. List of the specimens of dipterous insects in the collection of the British Museum London, V, I: 450-458, Tabanidae 450-545 pp.
- Yücel, Ş.** 1987. İç Anadolu Bölgesinde bulunan Tabanidae (Diptera) türleri üzerinde araştırmalar, Ankara Üni. Sağlık Bil. Ens. Doktora Tezi, 155 pp.