

**THE LONGHORNED BEETLES THAT ORIGINALLY  
DESCRIBED FROM WHOLE TERRITORIES OF TURKEY  
(COLEOPTERA: CERAMBYCOIDEA) PART II –  
CERAMBYCIDAE (CERAMBYCINAE AND STENOPTERINAE)**

**Hüseyin Özdikmen\***

\* Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 06500 Ankara, TURKEY. E-mail: ozdikmen@gazi.edu.tr

**[Özdikmen, H. 2012.** The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part II - Cerambycidae (Cerambycinae and Stenopterinae). *Munis Entomology & Zoology*, 7 (2): 699-707]

**ABSTRACT.** The paper is the second part of a planned serial work. It gives an annotated list of the Cerambycoidea [Cerambycidae (Cerambycinae and Stenopterinae)] that originally described from Turkey, with some important synonyms.

**KEY WORDS:** Cerambycinae, Stenopterinae, Cerambycidae, Cerambycoidea, Coleoptera, Turkey.

A series work is planned that is aim to expose the Turkish Cerambycoidea described from Turkey originally as possible as detailed and entirely by beginning from Vesperidae to Cerambycidae (Özdikmen, 2012a). The present study that includes the subfamilies Cerambycinae and Stenopterinae, is attempted as the second step of this aim.

Information in the present text is given in following order:

All taxa are accompanied by the author's name and description date. For each species, subspecies or important synonym, the original combination and type locality are provided under the taxon names. The status of endemism for Turkey is also indicated under these information.

In this paper, classification and nomenclature of the longhorn beetles suggested by Özdikmen (2012b) is followed chiefly. Within the genera and subgenera, the species are listed alphabetically.

Please see Löbl & Smetana (2010) for cited references in the text.

**SUPERFAMILY CERAMBYCOIDEA Latreille, 1802**

**FAMILY CERAMBYCIDAE Latreille, 1802: 211**

**SUBFAMILY CERAMBYCINAE Latreille, 1802: 211**

**TRIBE HESPEROPHANINI** Mulsant, 1839: 61

**SUBTRIBE HESPEROPHANINA** Mulsant, 1839: 61

**GENUS** *TRICHOFERUS* Wollaston, 1854: 427

**SUBGENUS -**

**SPECIES** *T. kotschyi* (Ganglbauer, 1883: 300)

**Orig. comb.:** *Hesperophanes kotschyi*

**Type loc.:** Asia Minor "Cilicia" (Turkey)

**SPECIES** *T. lunatus* (Szallies, 1994: 261)

**Orig. comb.:** *Hesperophanes lunatus*

**Type loc.:** Mardin: Hop pass (Turkey)

**SPECIES** *T. preissi* (Heyden, 1894: 85)

**Orig. comb.:** *Hesperophanes preissi*

**Type loc.:** Mardin (Turkey)

**Endemic species to Turkey.**

**SPECIES** *T. samai* Kadlec & Rejzek, 2001: 296

**Orig. comb.:** *Trichoferus samai*

**Type loc.:** Adiyaman: Nemrut Mt., Kahta env. (Turkey)

**Endemic species to Turkey.**

**SPECIES** *T. sbordonii* Sama, 1982: 217

**Orig. comb.:** *Trichoferus sbordonii*

**Type loc.:** Antalya: Alanya (Turkey)

**Endemic species to Turkey.**

**TRIBE CERAMBYCINI** Latreille, 1802: 211

**SUBTRIBE CERAMBYCINA** Latreille, 1802: 211

**GENUS CERAMBYX** Linnaeus, 1758: 388

**SUBGENUS CERAMBYX** Linnaeus, 1758: 388

**SPECIES** *C. heinzianus* Demelt, 1976: 65

**Orig. comb.:** *Cerambyx heinzianus*

**Type loc.:** Bingöl: Solhan (Turkey)

**Endemic species to Turkey.**

**SUBGENUS MICROCERAMBYX** Mikšić & Georgijevic, 1973: 22

**SPECIES** *C. scopolii* Fuessly, 1775: 12

**Orig. comb.:** *Cerambyx scopolii*

**Type loc.:** Switzerland

**SUBSPECIES** *C. s. nitidus* Pic, 1892: CXI

**Orig. comb.:** *Cerambyx scopolii* var. *nitidus*

**Type loc.:** "monts Amanus, pays d'Akbès" (Turkey: Hatay: Akbez)

**Endemic subspecies to Turkey.**

**TRIBE ROSALIINI** Fairmaire, 1864: 137

**SUBTRIBE** -

**GENUS ROSALIA** Audinet-Serville, 1834: 561

**SUBGENUS ROSALIA** Audinet-Serville, 1834: 561

**SPECIES** *R. alpina* (Linnaeus, 1758: 392)

**Orig. comb.:** *Cerambyx alpinus*

**Type loc.:** "Helvetia" (Switzerland)

**SUBSPECIES** *R. a. syriaca* Pic, 1895: CCLXXXV

**Orig. comb.:** *Rosalia alpina* var. *syriaca*

**Type loc.:** Hatay: Akbez (Turkey)

**Endemic subspecies to Turkey.**

**TRIBE TRACHYDERINI** Dupont, 1836: 1

**SUBTRIBE TRACHIDERINA** Dupont, 1836: 1

**GENUS PURPURICENUS** Dejean, 1821: 105

**SUBGENUS** -

**SPECIES** *P. bitlisiensis* Pic, 1902: 27

**Orig. comb.:** *Purpuricenus budensis* var. *bitlisiensis*

**Type loc.:** Bitlis (Turkey)

**Endemic species to Turkey.**

**SPECIES** *P. cornifrons* Sabbadini & Pesarini, 1992: 58

**Orig. comb.:** *Purpuricenus cornifrons*

**Type loc.:** Bingöl (Turkey)

**Endemic species to Turkey.**

**SPECIES** *P. interscapillatus* Plavilstshikov, 1937: 247 [RN]

**Orig. comb.:** *Purpuricenus interscapillatus*

**Type loc.:** Lebanon

**SUBSPECIES** *P. i. nudicollis* Demelt, 1968: 65

**Orig. comb.:** *Purpuricenus nudicollis*

**Type loc.:** İçel (Turkey)

**Endemic subspecies to Turkey.**

**SPECIES** *P. nigrotatatus* Pic, 1907: 169

**Orig. comb.:** *Purpuricenus budensis* var. *nigronotatus*

**Type loc.:** Adana: Karsanti (Turkey)

**Endemic species to Turkey.**

**SYNONYM**

*konradi* Bernhauer, 1976: 123

**Orig. comb.:** *Purpuricenus konradi*

**Type loc.:** Adana: Nurdağı pass (Turkey: Hatay prov.)

**GENUS** *CALCHAENESTHES* Kraatz, 1863: 97

**SUBGENUS -**

**SPECIES** *C. oblongomaculata* (Guerin-Meneville, 1844: 234)

**Orig. comb.:** *Callidium oblongomalatum*

**Type loc.:** Graecia (Greece)

**SYNONYM**

*nogeli* Frivaldszky, 1845: 183

**Orig. comb.:** *Callidium nogeli*

**Type loc.:** "Constantinopoly" (Turkey: İstanbul)

**TRIBE** CALLICHRMATINI Swainson & Shuckard, 1840: 293

**SUBTRIBE -**

**GENUS** *OSPHRANTERIA* Redtenbacher, 1850: 50

**SUBGENUS -**

**SPECIES** *O. coerulescens* Redtenbacher, 1850: 50

**Orig. comb.:** *Osphranteria coerulescens*

**Type loc.:** Shiraz (Iran)

**SUBSPECIES** *O. c. inaurata* Holzschuh, 1981: 98

**Orig. comb.:** *Osphranteria coerulescens inaurata*

**Type loc.:** Hakkari: Sat Mt.: Oramar (=Dağlıca) (Turkey)

**TRIBE** GRACILIINI Mulsant, 1839: 99

**SUBTRIBE -**

**GENUS** *HYBOMETOPIA* Ganglbauer, 1889: 282

**SUBGENUS -**

**SPECIES** *H. starcki* Ganglbauer, 1889: 285

**Orig. comb.:** *Hybometopia starcki*

**Type loc.:** Caucasus

**SUBSPECIES** *H. s. ivani* Sama, 1996: 106

**Orig. comb.:** *Hybometopia starcki ivani*

**Type loc.:** Kastamonu: Yaralıgöz (Turkey)

**TRIBE** OBRINI Mulsant, 1839: 95

**SUBTRIBE -**

**GENUS** *ANATOLOBRIUM* Adlbauer, 2004: 419

**SUBGENUS -**

**SPECIES** *A. eggeri* Adlbauer, 2004: 421

**Orig. comb.:** *Anatolobrium eggeri*

**Type loc.:** Antalya: Alanya (Turkey)

**Endemic species to Turkey.**

**TRIBE** DELLINI Fairmaire, 1864: 154

**SUBTRIBE -**

**GENUS** *DELAGRANGEUS* Pic, 1892: XCIII

**SUBGENUS** *DELAGRANGEUS* Pic, 1892: XCIII

**SPECIES** *D. angustissimus* Pic, 1892: XCIII

**Orig. comb.:** *Delagrangaeus angustissimus*

**Type loc.:** "Haute Syria" (Turkey: Hatay: Akbez)

**SYNONYM**

*asiaticus* Pic, 1920: 21

**Orig. comb.:** *Mimophymatodes asiaticus*

**Type loc.:** Konya: Akşehir (Turkey)

**SUBSPECIES** *D. a. angustissimus* Pic, 1892: XCIII  
**Orig. comb.:** *Delagrangeus angustissimus angustissimus*  
**Type loc.:** "Haute Syria" (Turkey: Hatay: Akbez)  
**Endemic subspecies to Turkey.**

**GENUS** *DEILUS* Audinet-Serville, 1834: 73

**SUBGENUS** -

**SPECIES** *D. rugosicollis* Rapuzzi & Sama, 2012: 668

**Orig. comb.:** *Deilus rugosicollis*

**Type loc.:** Lebanon, Syria, Turkey, Israel

**TRIBE** CALLIDIINI Kirby, 1837: 170

**SUBTRIBE** -

**GENUS** *ROPALOPUS* Mulsant, 1839: 40

**SUBGENUS** *ROPALOPUS* Mulsant, 1839: 40

**SPECIES** *R. hanae* Sama & Rejzek, 2002: 105

**Orig. comb.:** *Ropalopus hanae*

**Type loc.:** Muş: Buğlan pass (Turkey)

**Endemic species to Turkey.**

**SPECIES** *R. ledereri* (Fairmaire, 1866: 269)

**Orig. comb.:** *Rhopalopus ledereri*

**Type loc.:** Asie Mineure (Turkey: ?W Anatolia: ?İzmir)

**SUBSPECIES** *R. l. ledereri* (Fairmaire, 1866: 269)

**Orig. comb.:** *Rhopalopus ledereri*

**Type loc.:** Asie Mineure (Turkey: ?W Anatolia: ?İzmir)

**SUBSPECIES** *R. l. wittmeri* Demelt, 1970: 31

**Orig. comb.:** *Rhopalopus ledereri wittmeri*

**Type loc.:** Kahramanmaraş, Osmaniye (Turkey)

**SPECIES** *R. sculpturatus* (Pic, 1931: 9)

**Orig. comb.:** *Rhopalopus clavipes* var. *sculpturatus*

**Type loc.:** Tokat (Turkey)

**Endemic species to Turkey.**

**GENUS** *LEIODERES* Redtenbacher, 1849: 482

**SUBGENUS** -

**SPECIES** *L. tuerki* (Ganglbauer, 1886: 517)

**Orig. comb.:** *Callidium (Lioderes) türki*

**Type loc.:** Cilicia: Gülek (Bulgar-Dagh) (Turkey: İçel)

**GENUS** *CALLIDIUM* Fabricius, 1775: 187

**SUBGENUS** *CALLIDIUM* Fabricius, 1775: 187

**SPECIES** *C. syriacum* Pic, 1892: CXI

**Orig. comb.:** *Callidium ?aeneum* var. *syriacum*

**Type loc.:** "Nord Syrie" (Turkey: Hatay: Akbez not Syria)

**GENUS** *PHYMATODES* Mulsant, 1839: 47

**SUBGENUS** *MELASMETUS* Reitter, 1913: 39

**SPECIES** *P. femoralis* (Ménétriés, 1832: 228)

**Orig. comb.:** *Callidium femorale*

**Type loc.:** "Talyche" (Caucasus)

**SUBSPECIES** *P. f. demelti* Heyrovsky, 1962: 41

**Orig. comb.:** *Phymatodes femoralis demelti*

**Type loc.:** İstanbul (Turkey)

**Endemic subspecies to Turkey.**

**SUBGENUS** *PHYMATODELLUS* Reitter, 1913: 40

**SPECIES** *P. rufipes* (Fabricius, 1777: 232)

**Orig. comb.:** *Callidium rufipes*

**Type loc.:** "Germania" (Germany)

**SUBSPECIES** *P. r. syriacus* (Pic, 1891: 118)

**Orig. comb.:** *Callidium* (*Poecilium*) *rufipes* var. *syriacum*

**Type loc.:** "Syrie" (Turkey: ?Hatay: Akbez not Syria)

**SUBGENUS** *POECILIUM* Fairmaire, 1864: 134

**SPECIES** *P. kasnaki* (Sama, 2011: 826)

**Orig. comb.:** *Poecilium kasnaki*

**Type loc.:** Isparta: Yukarıgökdere (Turkey)

**Endemic species to Turkey.**

**SPECIES** *P. magnanii* (Sama & Rapuzzi, 1999: 468)

**Orig. comb.:** *Poecilium magnanii*

**Type loc.:** Antalya: Akseki: Çakılı pass (Turkey)

**Endemic species to Turkey.**

**TRIBE ANAGLYPTINI** Lacordaire, 1868: 404

**SUBTRIBE** -

**GENUS** *ANAGLYPTUS* Mulsant, 1839: 91

**SUBGENUS** *ANAGLYPTUS* Mulsant, 1839: 91

**SPECIES** *A. arabicus* (Küster, 1847: 95)

**Orig. comb.:** *Clytus arabicus*

**Type loc.:** Turkey

**SPECIES** *A. croceus* Pesarini & Sabbadini, 1997: 47

**Orig. comb.:** *Anaglyptus croceus*

**Type loc.:** İzmir: Bozdağ (Turkey)

**Endemic species to Turkey.**

**TRIBE CLYTINI** Mulsant, 1839: 70

**SUBTRIBE** -

**GENUS** *ISOTOMUS* Mulsant, 1862: 143

**SUBGENUS** -

**SPECIES** *I. comptus* (Mannerheim, 1825: 36)

**Orig. comb.:** *Clytus comptus*

**Type loc.:** "Imperii Rossici"

**SUBSPECIES** *I. c. meridionalis* Özdikmen & Aytar, 2012: 653

**Orig. comb.:** *Isotomus comptus meridionalis*

**Type loc.:** Osmaniye: Karatepe (Turkey)

**Endemic subspecies to Turkey.**

**SPECIES** *I. syriacus* (Pic, 1902: 25)

**Orig. comb.:** *Caloclytus syriacus*

**Type loc.:** "Syrie" (Turkey: Hatay not Syria)

**Endemic species to Turkey.**

**GENUS** *CHLOROPHORUS* Chevrolat, 186: 290

**SUBGENUS** *CRASSOFASCIATUS* Özdikmen, 2011: 538

**SPECIES** *C. convexifrons* Holzschuh, 1981: 100

**Orig. comb.:** *Chlorophorus convexifrons*

**Type loc.:** Samsun and Manisa (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. cursor* Rapuzzi & Sama, 1999: 331

**Orig. comb.:** *Chlorophorus cursor*

**Type loc.:** Bolu: Abant Lake (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. niehuisi* Adlbauer, 1992: 497

**Orig. comb.:** *Chlorophorus niehuisi*

**Type loc.:** Muş: Buğlan pass (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. oezdikmeni* Sama & Rapuzzi, 2011

**Orig. comb.:** *Chlorophorus oezdikmeni*

**Type loc.:** Kahramanmaraş: Andırın (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. robustior* (Pic, 1900: 11)  
**Orig. comb.:** *Clytanthus trifasciatus* var. *robustior*  
**Type loc.:** Konya (Turkey)  
**Endemic species to Turkey.**

**SUBGENUS** *PERDEROMACULATUS* Özdikmen, 2011: 537

**SPECIES** *C. graciosus* (Marseul, 1868: 203)  
**Orig. comb.:** *Clytus graciosus*  
**Type loc.:** Beyrouth env. (Lebanon)

**SUBSPECIES** *C. g. sparsus* (Marseul, 1868: 203)  
**Orig. comb.:** *Clytus (Clytanthus) sparsus*  
**Type loc.:** "Adalia, Lycischen Taurus" (Turkey: Antalya)  
**Endemic subspecies to Turkey.**

**SPECIES** *C. grosseri* Sama & Rapuzzi, 2011: 85  
**Orig. comb.:** *Chlorophorus grosseri*  
**Type loc.:** Şirnak: Mersindağı pass (Turkey)  
**Endemic species to Turkey.**

**SPECIES** *C. wewalkai* Holzschuh, 1969: 77  
**Orig. comb.:** *Chlorophorus wewalkai*  
**Type loc.:** Ankara: Kızılcahamam (Turkey)  
**Endemic species to Turkey.**

**SUBGENUS** *HUMEROMACULATUS* Özdikmen, 2011: 537

**SPECIES** *C. dinae* Rapuzzi & Sama, 1999: 329  
**Orig. comb.:** *Chlorophorus dinae*  
**Type loc.:** Hatay: Antakya: Şenköy (Turkey)

**SPECIES** *C. dominici* Sama, 1996: 110  
**Orig. comb.:** *Chlorophorus dominici*  
**Type loc.:** Kastamonu: Devrekani (Turkey)  
**Endemic species to Turkey.**

**SPECIES** *C. nivipictus* (Kraatz, 1879: 91)  
**Orig. comb.:** *Clytus nivipictus*  
**Type loc.:** "Cilicischen Taurus" (Turkey)

**GENUS** *TURANOCLYTUS* Sama, 1994: hevrolat, 1860: 456

**SUBGENUS** -

**SPECIES** *T. sieversi* (Ganglbauer, 1890: 429)  
**Orig. comb.:** *Clytus (Xylotrechus) sieversi*  
**Type loc.:** "Kasikoparan, (Armenia rossica)" (Turkey: Iğdır)

**SYNONYM**

*akbesianus* Pic, 1902: 17  
**Orig. comb.:** *Xylotrechus akbesianus*  
**Type loc.:** Hatay: Akbez (Turkey)

**GENUS** *PSEUDOSPHEGESTHES* Reitter, 1913: 50

**SUBGENUS** -

**SPECIES** *P. longitarsus* Holzschuh, 1974: 90  
**Orig. comb.:** *Pseudosphegthes longitarsus*  
**Type loc.:** Muş: Buğlan pass (Turkey)  
**Endemic species to Turkey.**

**SPECIES** *P. samai* Danilevsky, 2000: 44  
**Orig. comb.:** *Pseudosphegthes samai*  
**Type loc.:** Tokat: Niksar (Turkey)  
**Endemic species to Turkey.**

GENUS *CLYTUS* Laicharting, 1784: 88

SUBGENUS -

**SPECIES** *C. buglanicus* Kadlec, 2005: 106

**Orig. comb.:** *Clytus buglanicus*

**Type loc.:** Muş: Buğlan pass (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. ciliciensis* (Chevrolat, 1863: 334)

**Orig. comb.:** *Sphegesthes ciliciensis*

**Type loc.:** "Caramanie" (Turkey: Karaman)

**SPECIES** *C. gulekanus* Pic, 1904: 65

**Orig. comb.:** *Clytus gulekanus*

**Type loc.:** İçel: Gülek (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. insignitus* Fairmaire, 1866: 269 [DA]

**Orig. comb.:** *Clytus insignitus*

**Type loc.:** "Kisilgye-Aole" (Turkey: ?İzmir or Burdur: Kızılkaya)

**Endemic species to Turkey.**

**SPECIES** *C. kumalariensis* Johanides, 2001: 219

**Orig. comb.:** *Clytus kumalariensis*

**Type loc.:** Afyon: Kumalar Mt. (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. schneideri* Kiesenwetter, 1878: 313 [= 1879: 57]

**Orig. comb.:** *Clytus schneideri*

**Type loc.:** Caucasus

**SUBSPECIES** *C. s. inapicalis* Pic, 1895: 38

**Orig. comb.:** *Clytus inapicalis*

**Type loc.:** Erzurum (Turkey)

**Endemic subspecies to Turkey.**

**SPECIES** *C. schurmanni* Sama, 1996: 108

**Orig. comb.:** *Clytus schurmanni*

**Type loc.:** Ankara: Kızılcahamam (Turkey)

**Endemic species to Turkey.**

**SPECIES** *C. taurusiensis* (Pic, 1903: 139)

**Orig. comb.:** *Clytus (Clytanthus) taurusiensis*

**Type loc.:** "Taurus Mts." (Turkey)

**SUBFAMILY STENOPTERINAE Gistel, 1848: [9] (unnumbered section)**

**TRIBE STENOPTERINI Gistel, 1848: [9]**

**SUBTRIBE -**

GENUS *STENOPTERUS* Illiger, 1804: 120

SUBGENUS -

**SPECIES** *S. adlbaueri* Sama, 1995: 408

**Orig. comb.:** *Stenopterus adlbaueri*

**Type loc.:** Hakkari (Turkey)

**Endemic species to Turkey.**

**SPECIES** *S. atricornis* Pic, 1891: 102

**Orig. comb.:** *Stenopterus rufus* v. *atricornis*

**Type loc.:** "Syrie (Akbès)" (Turkey: Hatay: Akbez not Syria)

**SPECIES** *S. kraatzi* Pic, 1892: 21

**Orig. comb.:** *Stenopterus kraatzi*

**Type loc.:** "Asie Mineure, Smyrna" (Turkey: İzmir)

**Endemic species to Turkey.**

**SPECIES** *S. rufus* (Linnaeus, 1767: 642)

**Orig. comb.:** *Necydalis rufus*

**Type loc.:** "Europa australi" (Southern Europe)

**SUBSPECIES** *S. r. syriacus* Pic, 1892: 22

**Orig. comb.:** *Stenopterus rufus* v. *syriacus*

**Type loc.:** "Syrie (Akbès)" (Turkey: Hatay: Akbez not Syria)

**TRIBE MOLORCHINI** Gistel, 1848: [9] (unnumbered section)

**SUBTRIBE** -

**GENUS MOLORCHUS** Fabricius, 1792: 356

**SUBGENUS CAENOPTERA** Thomson, 1859: 150

**SPECIES** *M. abieticola* Holzschuh, 2007: 218

**Orig. comb.:** *Molorchus abieticola*

**Type loc.:** İçel: Namrun (Turkey)

**Endemic species to Turkey.**

**SPECIES** *M. juglandis* Sama, 1982: 219

**Orig. comb.:** *Molorchus juglandis*

**Type loc.:** Antalya: Alanya (Turkey)

**SUBGENUS MOLORCHUS** Fabricius, 1792: 356

**SPECIES** *M. kiesenwetteri* Mulsant & Rey, 1861: 189

**Orig. comb.:** *Molorchus kiesenwetteri*

**Type loc.:** not stated, probably Germany

**SUBSPECIES** *M. k. hircus* Abeille de Perrin, 1881: 133

**Orig. comb.:** *Molorchus hircus*

**Type loc.:** "Bloudan (Anti Liban)" (Southern Syria)

**SYNONYMS**

*anatolicus* Adlbauer, 1988: 275

**Orig. comb.:** *Molorchus kiesenwetteri anatolicus*

**Type loc.:** İçel: Erdemli (Turkey)

*angorensis* Pic, 1912: 3

**Orig. comb.:** *Caenoptera angorensis*

**Type loc.:** Ankara (Turkey)

*castanipennis* Reitter, 1912: 27

**Orig. comb.:** *Molorchus kiesenwetteri* ab. *castanipennis*

**Type loc.:** "Kleinasien, Araxestal" (Turkey)

**SPECIES** *M. malmusii* (Sama, 1995: 370)

**Orig. comb.:** *Glaphyra malmusii*

**Type loc.:** Bolu: Abant (Turkey)

**Endemic species to Turkey.**

**SPECIES** *M. marmottani* (Brisout, 1863: 118)

**Orig. comb.:** *Molorchus marmottani*

**Type loc.:** "Vernet les Bains" (France: Pyrenees)

**SUBSPECIES** *M. m. frischi* (Sama, 1995: 373)

**Orig. comb.:** *Glaphyra marmottani frischi*

**Type loc.:** Osmaniye: Zorkun plateau (Turkey)

**Endemic subspecies to Turkey.**

**SPECIES** *M. tenuitarsis* Holzschuh, 1981: 97

**Orig. comb.:** *Molorchus tenuitarsis*

**Type loc.:** İçel: Namrun (Turkey)

**Endemic species to Turkey.**

**SYNONYM**

*sterbai* Adlbauer, 1988: 277

**Orig. comb.:** *Molorchus sterbai*

**Type loc.:** Kahramanmaraş: Gökşun (Turkey)



**TRIBE BRACHYPTEROMINI** Sama, 2008: 229

**SUBTRIBE** -

**GENUS** *BRACHYPTEROMA* Heyden, 1863: 128 [NP]

**SUBGENUS** -

**SPECIES** *B. holtzi* Pic, 1905: 114

**Orig. comb.:** *Brachypteroma holtzi*

**Type loc.:** Taurus Mts. (Turkey)

**SPECIES** *B. ottomanum* Heyden, 1863: 128 [NP]

**Orig. comb.:** *Brachypteroma ottomanum*

**Type loc.:** European Turkey

**TRIBE HYBODERINI** Linsley, 1840: 367

**GENUS** *CALLIMUS* Mulsant, 1846: [5]

**SUBGENUS** *CALLIMUS* Mulsant, 1846: [5]

**SPECIES** *C. akbesianus* Pic, 1892: CXI

**Orig. comb.:** *Callimus akbesianus*

**Type loc.:** Hatay: Akbez (Turkey)

**Endemic species to Turkey.**

**SUBGENUS** *PROCALLIMUS* Pic, 1907: 7

**SPECIES** *C. egregius* Mulsant & Rey, 1863: 146

**Orig. comb.:** *Callimus egregius*

**Type loc.:** "La Caramanie" (Turkey: Karaman)

As a result of the present work, a total of 66 species group taxa (51 species and 15 subspecies) of Cerambycidae [Cerambycinae (40 sp. and 13 ssp.) and Stenopterinae (11 sp. and 2 ssp.)] are determined.

Among of them, 44 species group taxa (35 species and 9 subspecies) are endemic to Turkey now. So, endemism ratio of the longhorned beetles that originally described from whole territories of Turkey, is approximately 68 % [Cerambycidae [Cerambycinae (29 sp. and 8 ssp.) and Stenopterinae (6 sp. and 1 ssp.)]].

### ACKNOWLEDGEMENT

Special thanks to M. L. Danilevsky (Russia) for some information.

### LITERATURE CITED

**Löbl I. & Smetana A.** (ed.) 2010. Catalogue of Palaearctic Coleoptera, Vol. 6. Chrysomeloidea. Stenstrup: Apollo Books, 924 pp.

**Özdikmen, H.** 2012a. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part I - Vesperidae and Cerambycidae (Prioninae, Lepturinae, Necydalinae, Aseminae, Saphaninae and Dorcasominae). *Munis Entomology & Zoology*, 7 (1): 592-603.

**Özdikmen, H.** 2012b. Naked lists of Turkish Cerambycoidea and Chrysomeloidea (Coleoptera). *Munis Entomology & Zoology*, 7 (1): 51-108.