

**THE LONGHORNED BEETLES THAT ORIGINALLY
DESCRIBED FROM WHOLE TERRITORIES OF TURKEY
(COLEOPTERA: CERAMBYCOIDEA) PART III –
CERAMBYCIDAE (DORCADIONINAE)**

Hüseyin Özdikmen*

* Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 06500 Ankara, TURKEY. E-mail: ozdikmen@gazi.edu.tr

[Özdikmen, H. 2012. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part III - Cerambycidae (Dorcadioninae). Munis Entomology & Zoology, 7 (2): 759-779]

ABSTRACT. The paper is the third part of a planned serial work. It gives an annotated list of the Cerambycoidea [Cerambycidae (Dorcadioninae)] that originally described from Turkey, with some important synonyms.

KEY WORDS: Dorcadioninae, Cerambycidae, Cerambycoidea, Coleoptera, Turkey.

A series work is planned that is aim to expose the Turkish Cerambycoidea described from Turkey originally as possible as detailed and entirely by beginning from Vesperidae to Cerambycidae (Özdikmen, 2012a,b). The present study that includes the subfamily Dorcadioninae, is attempted as the third step of this aim. Although, The Turkish Dorcadioninae was evaluated by Özdikmen (2010) in detailed.

Information in the present text is given in following order:

All taxa are accompanied by the author's name and description date. For each species, subspecies or important synonym, the original combination and type locality are provided under the taxon names. The status of endemism for Turkey is also indicated under these information.

In this paper, classification and nomenclature of the longhorn beetles suggested by Özdikmen (2012c) is followed chiefly. Within the genera and subgenera, the species are listed alphabetically.

Please see Löbl & Smetana (2010) for cited references in the text.

SUPERFAMILY CERAMBYCOIDEA Latreille, 1802

FAMILY CERAMBYCIDAE Latreille, 1802: 211

SUBFAMILY DORCADIONINAE Swainson, 1840: 290

TRIBE DORCADIONINI Swainson, 1840: 290

SUBTRIBE -

GENUS DORCADION Dalman, 1817: 397

SUBGENUS CARINATODORCADION Breuning, 1943: 524

SPECIES *D. ingeae* Peks, 1993: 3

Orig. comb.: *Dorcadion ingeae*

Type loc.: Edirne: Keşan (Turkey)

Endemic species to Turkey.

SUBGENUS CRIBRIDORCADION Pic, 1901: 12

SPECIES *D. abstersum* Holzschuh, 1982: 75

Orig. comb.: *Dorcadion abstersum*

Type loc.: Hakkari: Sat Mt. (Turkey)

Endemic species to Turkey.

SPECIES *D. accola* Heyden, 1894: 87
Orig. comb.: *Dorcadion accola*
Type loc.: Mardin (Turkey)
Endemic species to Turkey.

SYNONYM

mardinense Pic, 1900: 12
Orig. comb.: *Dorcadion accola* v. *mardinense*
Type loc.: Mardin (Turkey)

SPECIES *D. afflictum* Pesarini & Sabbadini, 1999: 54
Orig. comb.: *Dorcadion afflictum*
Type loc.: Çorum (Turkey)
Endemic species to Turkey.

SPECIES *D. akpinarense* Bernhauer & Peks, 2010
Orig. comb.: *Dorcadion akpinarense*
Type loc.: İçel: Anamur (Turkey)
Endemic species to Turkey.

SPECIES *D. albicolle* Breuning, 1943: 89
Orig. comb.: *Dorcadion albicolle*
Type loc.: Konya (Turkey)
Endemic species to Turkey.

SPECIES *D. albolineatum* Küster, 1847: 86
Orig. comb.: *Dorcadion albolineatum*
Type loc.: İstanbul (Turkey)
Endemic species to Turkey.

SYNONYM

byzantinum Thomson, 1867: 69
Orig. comb.: *Dorcadion byzantinum*
Type loc.: Mardin (Turkey)
Endemic species to Turkey.

SPECIES *D. albonotatum* Pic, 1895: 39
Orig. comb.: *Dorcadion albonotatum*
Type loc.: Bitlis: Sipan Mt. (Turkey)
Endemic species to Turkey.

SPECIES *D. amanense* Breuning, 1943: 94
Orig. comb.: *Dorcadion amanense*
Type loc.: Amanos Mts. (Turkey)
Endemic species to Turkey.

SPECIES *D. anaticum* Pic, 1900: 12
Orig. comb.: *Dorcadion anaticum*
Type loc.: Konya (Turkey)
Endemic species to Turkey.

SYNONYM

rosinae Daniel in Bodemeyer, 1900: 169
Orig. comb.: *Dorcadion rosinae*
Type loc.: Konya: Akşehir: Sultan Mt. (Turkey)

SPECIES *D. arcivagum* Thomson, 1867: 121
Orig. comb.: *Dorcadion arcivagum*
Type loc.: İçel: Gülek, Taurus Mts. (Turkey)
Endemic species to Turkey.

SYNONYM

cilicium Pic, 1904: 4
Orig. comb.: *Dorcadion arcivagum* var. *cilicium*
Type loc.: Cilicia (Turkey)

SPECIES *D. arдахense* Breuning, 1975: 10

Orig. comb.: *Dorcadion arдахense*

Type loc.: Ardahan: Yalnızçam Mts. (Turkey)

Endemic species to Turkey.

SPECIES *D. bangi* Heyden, 1894: 89

Orig. comb.: *Dorcadion bangi*

Type loc.: Kastamonu (Turkey)

Endemic species to Turkey.

SUBSPECIES *D. b. bangi* Heyden, 1894: 89

Orig. comb.: *Dorcadion bangi bangi*

Type loc.: Kastamonu (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. b. heinzorum* Braun, 1975: 17

Orig. comb.: *Dorcadion bang heinzorum*

Type loc.: Kastamonu, Çorum (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. b. roridum* Pesarini & Sabbadini, 1999: 55

Orig. comb.: *Dorcadion bandi roridum*

Type loc.: Bolu: Gerede (Turkey)

Endemic subspecies to Turkey.

SPECIES *D. banjkovski* Plavilstshikov, 1958: 147

Orig. comb.: *Dorcadion banjkovski*

Type loc.: Erzurum (Turkey)

Endemic species to Turkey.

SPECIES *D. beloni* Pic, 1891: LXXVII

Orig. comb.: *Dorcadion beloni*

Type loc.: Siberia? (Turkey not Siberia)

Endemic species to Turkey.

SPECIES *D. bernhauerorum* Peks, 2010

Orig. comb.: *Dorcadion bernhauerorum*

Type loc.: Bursa: İznik (Turkey)

Endemic species to Turkey.

SPECIES *D. bisignatum* Jakovlev, 1900: 66

Orig. comb.: *Dorcadion bisignatum*

Type loc.: Artvin (Turkey)

Endemic species to Turkey.

SYNONYM

artvinicum Plavilstshikov, 1958: 4

Orig. comb.: *Dorcadion bisignatum* ab. *artvinicum*

Type loc.: Artvin (Turkey)

SPECIES *D. bithyniense* Chevrolat, 1856: 88

Orig. comb.: *Dorcadion bithyniense*

Type loc.: Bursa (Turkey)

Endemic species to Turkey.

SPECIES *D. blandulum* Holzschuh, 1977: 131

Orig. comb.: *Dorcadion blandulus*

Type loc.: Bingöl: Kuruca pass (Turkey)

Endemic species to Turkey.

SPECIES *D. bodemeyeri* K. Daniel, 1900: 140

Orig. comb.: *Dorcadion bodemeyeri*

Type loc.: "Asia Minor" (Turkey: Phrygia and Lycaonia)

Endemic species to Turkey.

- SPECIES** *D. boluense* Breuning, 1962: 38
Orig. comb.: *Dorcadion boluense*
Type loc.: Bolu: Bolu Mts. (Turkey)
Endemic species to Turkey.
- SUBSPECIES** *D. b. boluense* Breuning, 1962: 38
Orig. comb.: *Dorcadion boluense boluense*
Type loc.: Bolu: Bolu Mts. (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. b. corallinum* Pesarini & Sabbadini, 1999: 53
Orig. comb.: *Dorcadion boluense corallinum*
Type loc.: Ankara: Kızılcahamam, Çamlıdere (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. b. imitator* Pesarini & Sabbadini, 1999: 53
Orig. comb.: *Dorcadion boluense imitator*
Type loc.: Ankara: Karagöl (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. boszdagense* Fairmaire, 1866: 275
Orig. comb.: *Dorcadion boszdagense*
Type loc.: İzmir: Bozdağ (Turkey)
Endemic species to Turkey.
- SPECIES** *D. bouilloni* Breuning & Ruspoli, 1975: 116
Orig. comb.: *Dorcadion bouilloni*
Type loc.: Ordu: Mesudiye (Turkey)
Endemic species to Turkey.
- SPECIES** *D. brauni* Breuning, 1979: 92
Orig. comb.: *Dorcadion brauni*
Type loc.: Denizli: Tavas valley (Turkey)
Endemic species to Turkey.
- SPECIES** *D. bremeri* Breuning, 1981: 181
Orig. comb.: *Dorcadion bremeri*
Type loc.: Afyon: Akyarma (Turkey)
Endemic species to Turkey.
- SPECIES** *D. bulgharmaadense* Breuning, 1946: 116
Orig. comb.: *Dorcadion bulgharmaadense*
Type loc.: Niğde: Ulukışla (Maden village) (Turkey)
Endemic species to Turkey.
- SPECIES** *D. cachinno* Thomson, 1867: 120
Orig. comb.: *Dorcadion cachinno*
Type loc.: "Caramanie" (Turkey: Karaman)
Endemic species to Turkey.
- SYNONYM**
immersum Tournier, 1872: 281
Orig. comb.: *Dorcadion immersum*
Type loc.: Turkey
- SPECIES** *D. carinipenne* Pic, 1900: 13
Orig. comb.: *Dorcadion carinipenne*
Type loc.: Hatay: Akbez (Turkey not Syria)
Endemic species to Turkey.
- SPECIES** *D. carolisturani* Breuning & Ruspoli, 1971: 128
Orig. comb.: *Dorcadion carolisturani*
Type loc.: Giresun: Kümbet plateau (Turkey)
Endemic species to Turkey.

- SPECIES** *D. cinctellum* Fairmaire, 1866: 272
Orig. comb.: *Dorcadion cinctellum*
Type loc.: "Kisilgye-Aole" (Turkey: ?İzmir or Burdur: Kızılkaya)
Endemic species to Turkey.
- SPECIES** *D. coiffaiti* Breuning, 1962: 392
Orig. comb.: *Dorcadion coiffaiti*
Type loc.: Balıkesir: Manyas (Turkey)
Endemic species to Turkey.
- SPECIES** *D. condensatum* Küster, 1852: 92
Orig. comb.: *Dorcadion condensatum*
Type loc.: NW Turkey
Endemic species to Turkey.
- SPECIES** *D. confluens* Fairmaire, 1866: 274
Orig. comb.: *Dorcadion confluens*
Type loc.: İzmir: Bozdağ (Turkey)
Endemic species to Turkey.
- SPECIES** *D. crux* (Bilberg, 1817: 175)
Orig. comb.: *Lamia crux*
Type loc.: "Smyrna" (Turkey: İzmir)
Endemic species to Turkey.
- SPECIES** *D. culminicola* Thomson, 1867: 122
Orig. comb.: *Dorcadion culminicola*
Type loc.: "Caramania" (Turkey: Taurus Mts.)
Endemic species to Turkey.
- SPECIES** *D. delagrangei* Pic, 1894: 110
Orig. comb.: *Dorcadion delagrangei*
Type loc.: Hatay: Akbez (Turkey)
Endemic species to Turkey.
- SYNONYMS**
ardinense Pic, 1900: 12
Orig. comb.: *Dorcadion accola* var. *ardinense*
Type loc.: Mardin (Turkey)
- fuscopubescens* Pic, 1900: 12
Orig. comb.: *Dorcadion delagrangei* var. *fuscopubescens*
Type loc.: SE Turkey
- SPECIES** *D. deyrollei* Ganglbauer, 1884: 482
Orig. comb.: *Dorcadion deyrollei*
Type loc.: Bitlis (Turkey)
Endemic species to Turkey.
- SPECIES** *D. dimidiatum* Motschulsky, 1838: 186
Orig. comb.: *Dorcadion deyrollei*
Type loc.: Erivan (Armenia)
- SUBSPECIES** *D. d. korgei* Breuning, 1966: 21
Orig. comb.: *Dorcadion dimidiatum korgei*
Type loc.: Tunceli: Pülümür (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. d. kelkiticum* Özdikmen & Hasbenli, 2004: 23
Orig. comb.: *Dorcadion dimidiatum kelkiticum*
Type loc.: Gümüşhane: Kelkit (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. divisum* Germar, 1839: 15
Orig. comb.: *Dorcadion divisum*
Type loc.: "Smyrna" (Turkey: İzmir)

- SUBSPECIES** *D. d. divisum* Germar, 1839: 15
Orig. comb.: *Dorcadion divisum*
Type loc.: "Smyrna" (Turkey: İzmir)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. d. dissimile* Ganglbauer, 1884: 458
Orig. comb.: *Dorcadion divisum* var. *dissimile*
Type loc.: European Turkey
Endemic subspecies to Turkey.
- SUBSPECIES** *D. d. intercisum* Kraatz, 1873: 66
Orig. comb.: *Dorcadion divisum* var. *intercisum*
Type loc.: Sivas, Cilicia, Kurdistan (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. d. loratum* Thomson, 1867: 123
Orig. comb.: *Dorcadion loratum*
Type loc.: "Smyrna" (Turkey: İzmir)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. d. oedemischense* Heyrovsky, 1932: 104
Orig. comb.: *Dorcadion divisum* var. *oedemischense*
Type loc.: İzmir: Ödemiş (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. d. subdivisum* Breuning, 1955: 263
Orig. comb.: *Dorcadion divisum subdivisum*
Type loc.: Ankara (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. dobroljanskii* Suvorov, 1915: 116
Orig. comb.: *Dorcadion dobroljanskii*
Type loc.: "Erivansk. g." (Turkey)
Endemic species to Turkey.
- SPECIES** *D. drusoides* Breuning, 1962: 375
Orig. comb.: *Dorcadion drusoides*
Type loc.: Amanos Mts. (Turkey not Syria)
- SPECIES** *D. elazigi* Fuchs & Breuning, 1971: 439
Orig. comb.: *Dorcadion elazigi*
Type loc.: Elazığ and Muş: Buğlan pass (Turkey)
Endemic species to Turkey.
- SPECIES** *D. enricisturanii* Breuning & Ruspoli, 1971: 127
Orig. comb.: *Dorcadion enricisturanii*
Type loc.: Kastamonu: Tosya (Turkey)
Endemic species to Turkey.
- SUBSPECIES** *D. e. enricisturanii* Breuning & Ruspoli, 1971: 127
Orig. comb.: *Dorcadion enricisturanii enricisturanii*
Type loc.: Kastamonu: Tosya (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. e. densepunctatum* Braun, 1978: 107
Orig. comb.: *Dorcadion enricisturanii densepunctatum*
Type loc.: Samsun: Vezirköprü (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. equestre* (Laxmann, 1770: 596)
Orig. comb.: *Cerambyx equestris*
Type loc.: S Russia
- SUBSPECIES** *D. e. nogelli* Fairmaire, 1866: 270
Orig. comb.: *Dorcadion nogelli*
Type loc.: İzmir: Bozdağ (Turkey)
Endemic subspecies to Turkey.

SPECIES *D. ferruginipes* Ménetries, 1836: 151

Orig. comb.: *Dorcadion ferruginipes*

Type loc.: İstanbul (Turkey)

SYNONYM

thoracicum Küster, 1846: 90

Orig. comb.: *Dorcadion thoracicum*

Type loc.: İstanbul (Turkey)

SPECIES *D. formosum* Kraatz, 1871: 411

Orig. comb.: *Dorcadion formosum*

Type loc.: not stated

Endemic species to Turkey.

SUBSPECIES *D. f. formosum* Kraatz, 1871: 411

Orig. comb.: *Dorcadion formosum formosum*

Type loc.: not stated

Endemic subspecies to Turkey.

SUBSPECIES *D. f. ponticum* Breuning, 1964: 1

Orig. comb.: *Dorcadion formosum ponticum*

Type loc.: "Pontic Alpes" (Turkey)

Endemic subspecies to Turkey.

SPECIES *D. gallipolitanum* Thomson, 1867: 59

Orig. comb.: *Dorcadion gallipolitanum*

Type loc.: "Gallipoli" (Turkey: Çanakkale)

SUBSPECIES *D. g. gallipolitanum* Thomson, 1867: 59

Orig. comb.: *Dorcadion gallipolitanum gallipolitanum*

Type loc.: "Gallipoli" (Turkey: Çanakkale)

SYNONYMS

suturaalba Desbrochers, 1870: 125

Orig. comb.: *Dorcadion suturaalba*

Type loc.: "Gallipoli" (Turkey: Çanakkale)

gandolphei Tournier, 1872: 288

Orig. comb.: *Dorcadion gandolphei*

Type loc.: "Gallipoli" (Turkey: Çanakkale)

SPECIES *D. gebzeense* Breuning, 1974: 148

Orig. comb.: *Dorcadion gebzeense*

Type loc.: Kocaeli: İzmit; Gebze (Turkey)

Endemic species to Turkey.

SPECIES *D. glabricolle* Breuning, 1943: 92

Orig. comb.: *Dorcadion glabricolle*

Type loc.: Turkey (Anatolia and Kurdistan)

Endemic species to Turkey.

SPECIES *D. halepense* Kraatz, 1873: 72

Orig. comb.: *Dorcadion halepense*

Type loc.: Aleppo (Syria)

SUBSPECIES *D. h. sehitekamilense* Özdikmen et al., 2012: 595

Orig. comb.: *Dorcadion halepense sehitekamilense*

Type loc.: Gaziantep: Şehitkamil (Turkey)

Endemic subspecies to Turkey.

SPECIES *D. hampii* Mulsant & Rey, 1863: 157

Orig. comb.: *Dorcadion hampii*

Type loc.: ["La Perse"] clearly mislabeled, should be Turkey

Endemic species to Turkey.

- SUBSPECIES** *D. h. hampii* Mulsant & Rey, 1863: 157
Orig. comb.: *Dorcadion hampii hampii*
Type loc.: ["La Perse"] clearly mislabeled, should be Turkey
Endemic species to Turkey.
- SUBSPECIES** *D. h. aureovittatum* Kraatz, 1873: 81
Orig. comb.: *Dorcadion hampii aureovittatum*
Type loc.: Sivas (Turkey)
Endemic species to Turkey.
- SPECIES** *D. heinzi* Breuning, 1964: 1
Orig. comb.: *Dorcadion heinzi*
Type loc.: Giresun: Eğribel pass (Turkey)
Endemic species to Turkey.
- SPECIES** *D. holtzi* Pic, 1905: 115
Orig. comb.: *Dorcadion culminicola* var. *holtzi*
Type loc.: Cilician Taurus: Dümbelek Mt. (Turkey)
Endemic species to Turkey.
- SPECIES** *D. holzschuhi* Breuning, 1974: 148
Orig. comb.: *Dorcadion holzschuhi*
Type loc.: Muş: Buğlan pass (Turkey)
Endemic species to Turkey.
- SPECIES** *D. iconiense* Daniel, 1900: 140
Orig. comb.: *Dorcadion iconiense*
Type loc.: Konya (Turkey)
Endemic species to Turkey.
- SPECIES** *D. infernale* Mulsant & Rey, 1863: 158
Orig. comb.: *Dorcadion infernale*
Type loc.: ["La Perse"] clearly mislabeled, should be Turkey
Endemic species to Turkey.
- SUBSPECIES** *D. i. infernale* Mulsant & Rey, 1863: 158
Orig. comb.: *Dorcadion infernale infernale*
Type loc.: ["La Perse"] clearly mislabeled, should be Turkey
Endemic subspecies to Turkey.
- SUBSPECIES** *D. i. asperatum* Breuning, 1947: 169
Orig. comb.: *Dorcadion infernale asperatum*
Type loc.: Diyarbakır (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. i. edremitense* Breuning, 1966: 20
Orig. comb.: *Dorcadion infernale edremitense*
Type loc.: Balıkesir: Edremit (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. inikliense* Bernhauer & Peks, 2010
Orig. comb.: *Dorcadion inikliense*
Type loc.: Bursa: İznik, near lake (Turkey)
Endemic species to Turkey.
- SPECIES** *D. inspersum* Holzschuh, 1982: 76
Orig. comb.: *Dorcadion inspersum*
Type loc.: Van: Kuzgunkıran (Turkey)
Endemic species to Turkey.
- SPECIES** *D. intrusum* Pesarini & Sabbadini, 2009
Orig. comb.: *Dorcadion intrusum*
Type loc.: Edirne: near Gelibolu (Turkey)
Endemic species to Turkey.

SPECIES *D. investitum* Breuning, 1970 in Braun, 1978: 111
Orig. comb.: *Dorcadion investitum*
Type loc.: Erzurum: Horasan: Tahir pass (Turkey)
Endemic species to Turkey.

SPECIES *D. ispartense* Breuning, 1962: 394
Orig. comb.: *Dorcadion ispartense*
Type loc.: Isparta: Bozanönü (Turkey)
Endemic species to Turkey.

SPECIES *D. ivani* Pesarini & Sabbadini, 2011
Orig. comb.: *Dorcadion ivani*
Type loc.: Bingöl (Turkey)
Endemic species to Turkey.

SPECIES *D. jacovleviellum* Plavilstshikov, 1951: 120
Orig. comb.: *Dorcadion jacovleviellum*
Type loc.: Erzurum (Turkey)
Endemic species to Turkey.

SPECIES *D. janatai* Kadlec, 2006: 11
Orig. comb.: *Dorcadion janatai*
Type loc.: Tunceli: Karaoğlan Mt. (Turkey)
Endemic species to Turkey.

SPECIES *D. kagyzmanicum* Suvorov, 1915: 120
Orig. comb.: *Dorcadion kagyzmanicum*
Type loc.: Kars: Kağızman (Turkey)
Endemic species to Turkey.

SPECIES *D. karsense* Suvorov, 1915: 118
Orig. comb.: *Dorcadion karsense*
Type loc.: Kars: Kağızman (Turkey)
Endemic species to Turkey.

SPECIES *D. kasikoporanum* Pic, 1902: 10
Orig. comb.: *Dorcadion kasikoporanum*
Type loc.: Iğdır: Tuzluca: Kazkoparan village (Turkey)
Endemic species to Turkey.

SPECIES *D. kazanciense* Bernhauer & Peks, 2010
Orig. comb.: *Dorcadion kazanciense*
Type loc.: İçel: Anamur (Turkey)
Endemic species to Turkey.

SPECIES *D. kindermanni* Walth, 1838: 470
Orig. comb.: *Dorcadion kindermanni*
Type loc.: Turkey
Endemic species to Turkey.

SYNONYM
griseum Walth, 1838: 470
Orig. comb.: *Dorcadion griseum*
Type loc.: Turkey

SPECIES *D. kraetschmeri* Bernhauer, 1988: 98
Orig. comb.: *Dorcadion kraetschmeri*
Type loc.: Antalya: Akseki (Turkey)
Endemic species to Turkey.

SPECIES *D. kurdistanum* Breuning, 1944: 12
Orig. comb.: *Dorcadion kurdistanum*
Type loc.: Diyarbakır (Turkey)
Endemic species to Turkey.

SPECIES *D. kurucanum* Holzschuh, 2007: 249

Orig. comb.: *Dorcadion kurucanum*

Type loc.: Elazığ: Kuruca pass (Turkey)

Endemic species to Turkey.

SPECIES *D. ladikanum* Braun, 1976: 173

Orig. comb.: *Dorcadion ladikanum*

Type loc.: Samsun: Ladik (Turkey)

Endemic species to Turkey.

SYNONYM

ladikense Breuning, 1975: 10 [HN]

Orig. comb.: *Dorcadion ladikense*

Type loc.: Samsun: Ladik (Turkey)

SPECIES *D. laeve* Faldermann, 1837: 278

Orig. comb.: *Dorcadion laeve*

Type loc.: Iran

SUBSPECIES *D. l. micula* Plavilstshikov, 1937: 26

Orig. comb.: *Dorcadion micula*

Type loc.: Erzurum: Oltu (Turkey)

Endemic subspecies to Turkey.

SPECIES *D. lameerei* Théry, 1896: 109

Orig. comb.: *Dorcadion lameerei*

Type loc.: Kastamonu (Turkey)

Endemic species to Turkey.

SPECIES *D. ledouxi* Breuning, 1974: 65

Orig. comb.: *Dorcadion ledouxi*

Type loc.: Erzurum: Ilıca (Turkey)

Endemic species to Turkey.

SPECIES *D. linderi* Tournier, 1872: 285

Orig. comb.: *Dorcadion linderi*

Type loc.: ["Espagne"] clearly mislabeled, should be Turkey

Endemic species to Turkey.

SYNONYMS

kollari Kraatz, 1873: 87

Orig. comb.: *Dorcadion kollari*

Type loc.: Amasya (Turkey)

henrici Breuning, 1975: 10 [HN]

Orig. comb.: *Dorcadion henrici*

Type loc.: Tokat (Turkey)

SPECIES *D. lodosi* Sabbadini & Pesarini, 1992: 29

Orig. comb.: *Dorcadion lodosi*

Type loc.: Karaman: Bucakkaşla (Turkey)

Endemic species to Turkey.

SUBSPECIES *D. l. lodosi* Sabbadini & Pesarini, 1992: 29

Orig. comb.: *Dorcadion lodosi*

Type loc.: Karaman: Bucakkaşla (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. l. abanozense* Bernhauer & Peks, 2010

Orig. comb.: *Dorcadion lodosi abanozense*

Type loc.: İçel: Anamur (Turkey)

Endemic subspecies to Turkey.

SPECIES *D. lohsei* Braun, 1976: 254

Orig. comb.: *Dorcadion lohsei*

Type loc.: İçel: Sertavul pass (Turkey)

Endemic species to Turkey.

SPECIES *D. longulum* Breuning, 1943: 89

Orig. comb.: *Dorcadion longulum*

Type loc.: İzmir: Bozdağ (Turkey)

Endemic species to Turkey.

SPECIES *D. maceki* Holzschuh, 1965: 41

Orig. comb.: *Dorcadion maceki*

Type loc.: Bolu: Abant (Turkey)

Endemic species to Turkey.

SPECIES *D. martini* Bernhauer, 1988: 100

Orig. comb.: *Dorcadion martini*

Type loc.: Ankara: Nallıhan (Turkey)

Endemic species to Turkey.

SPECIES *D. menradi* Holzschuh, 1989: 172

Orig. comb.: *Dorcadion menradi*

Type loc.: Kahramanmaraş: Afşin (Turkey)

Endemic species to Turkey.

SPECIES *D. merkli* Ganglbauer, 1884: 506

Orig. comb.: *Dorcadion merkli*

Type loc.: Akdağ, ?Kütahya (Turkey)

Endemic species to Turkey.

SPECIES *D. micans* Thomson, 1867: 61

Orig. comb.: *Dorcadion micans*

Type loc.: "Armenia" (clearly N Turkey)

Endemic species to Turkey.

SUBSPECIES *D. m. micans* Thomson, 1867: 61

Orig. comb.: *Dorcadion micans micans*

Type loc.: "Armenia" (clearly N Turkey)

Endemic species to Turkey.

SYNONYMS

sericatum Kraatz, 1873: 98

Orig. comb.: *Dorcadion sericatum*

Type loc.: Ankara, Bursa (Turkey)

macropus Kraatz, 1873: 99

Orig. comb.: *Dorcadion macropus*

Type loc.: Amasya, Tokat (Turkey)

SUBSPECIES *D. m. susheriense* Breuning, 1970: 97

Orig. comb.: *Dorcadion micans susheriense*

Type loc.: Sivas: Susehri (Turkey)

Endemic species to Turkey.

SPECIES *D. miminfernale* Breuning, 1970: 98

Orig. comb.: *Dorcadion miminfernale*

Type loc.: Kahramanmaraş: Binboğa Mts. (Turkey)

Endemic species to Turkey.

SPECIES *D. muchei* Breuning, 1962: 38

Orig. comb.: *Dorcadion muchei*

Type loc.: Çankırı (Turkey)

Endemic species to Turkey.

SPECIES *D. multimaculatum* Pic, 1932: 22

Orig. comb.: *Dorcadion multimaculatum*

Type loc.: İzmir: Ödemiş (Turkey)

Endemic species to Turkey.

- SPECIES** *D. narlianum* Özdikmen, Mercan & Cihan, 2012: 550
Orig. comb.: *Dorcadion narlianum*
Type loc.: Kahramanmaraş: Narlı (Turkey)
Endemic species to Turkey.
- SPECIES** *D. nigrostriatum* Adlbauer, 1982: 104
Orig. comb.: *Dorcadion nigrostriatum*
Type loc.: Tunceli (Turkey)
Endemic species to Turkey.
- SPECIES** *D. nihalae* Rapuzzi & Sama, 2012: 671
Orig. comb.: *Dorcadion nihalae*
Type loc.: Sivas: Zara (Turkey)
Endemic species to Turkey.
- SPECIES** *D. nobile* Hampe, 1852: 313
Orig. comb.: *Dorcadion nobile*
Type loc.: not stated
Endemic species to Turkey.
- SPECIES** *D. obsoletum* Kraatz, 1873: 78
Orig. comb.: *Dorcadion obsoletum*
Type loc.: European Turkey
- SPECIES** *D. obtusum* Breuning, 1944: 13
Orig. comb.: *Dorcadion obtusum*
Type loc.: Turkey
Endemic species to Turkey.
- SUBSPECIES** *D. o. obtusum* Breuning, 1944: 13
Orig. comb.: *Dorcadion obtusum obtusum*
Type loc.: Turkey
Endemic subspecies to Turkey.
- SUBSPECIES** *D. o. marashense* Breuning, 1948: 59
Orig. comb.: *Dorcadion obtusum marashense*
Type loc.: Kahramanmaraş (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. oezdurali* Önalp, 1988: 361
Orig. comb.: *Dorcadion oezdurali*
Type loc.: Kahramanmaraş: Ahrır Mt. (Turkey)
Endemic species to Turkey.
- SPECIES** *D. olympicum* Kraatz, 1873: 78
Orig. comb.: *Dorcadion olympicum*
Type loc.: Bursa (Turkey)
Endemic species to Turkey.
- SUBSPECIES** *D. o. olympicum* Kraatz, 1873: 78
Orig. comb.: *Dorcadion olympicum olympicum*
Type loc.: Bursa (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. o. convexum* Breuning, 1943: 90
Orig. comb.: *Dorcadion convexum*
Type loc.: Akdağ, ?Kütahya (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. o. flavosuturale* Krätschmer, 1987: 340
Orig. comb.: *Dorcadion olympicum flavosuturale*
Type loc.: Bursa: Uluabat Lake (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. ortrudae* Braun, 1978: 185
Orig. comb.: *Dorcadion ortrudae*
Type loc.: Kahramanmaraş: Göksun (Turkey)
Endemic species to Turkey.

SPECIES *D. paracinerarium* Breuning, 1974: 149

Orig. comb.: *Dorcadion paracinerarium*

Type loc.: İçel: Mut: Sertavul pass (Turkey)

Endemic species to Turkey.

SPECIES *D. pararufipenne* Braun, 1976: 257

Orig. comb.: *Dorcadion pararufipenne*

Type loc.: Bolu: Abant (Turkey)

Endemic species to Turkey.

SUBSPECIES *D. p. pararufipenne* Braun, 1976: 257

Orig. comb.: *Dorcadion pararufipenne pararufipenne*

Type loc.: Bolu: Abant (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. p. rassei* Braun, 1976: 261

Orig. comb.: *Dorcadion pararufipenne rassei*

Type loc.: Bolu: Gerede and central (Turkey)

Endemic subspecies to Turkey.

SPECIES *D. parescherichi* Breuning, 1966: 146

Orig. comb.: *Dorcadion parescherichi*

Type loc.: Çorum: Boğazkale (Turkey)

Endemic species to Turkey.

SPECIES *D. pavesii* Pesarini & Sabbadini, 1999: 57

Orig. comb.: *Dorcadion pavesii*

Type loc.: Hakkari: Yüksekova (Turkey)

Endemic species to Turkey.

SPECIES *D. peksi* Bernhauer, 2010

Orig. comb.: *Dorcadion peksi*

Type loc.: İçel: Anamur (Turkey)

Endemic species to Turkey.

SPECIES *D. pergamenum* Pesarini & Sabbadini, 2009

Orig. comb.: *Dorcadion pergamenum*

Type loc.: İzmir: Bergama (Turkey)

Endemic species to Turkey.

SPECIES *D. petrovitzi* Heyrovsky, 1964: 97

Orig. comb.: *Dorcadion petrovitzi*

Type loc.: İçel (Turkey)

Endemic species to Turkey.

SPECIES *D. piochardi* Kraatz, 1873: 85

Orig. comb.: *Dorcadion piochardi*

Type loc.: Samsun, Amasya, Akdağ (?Kütahya) (Turkey)

Endemic species to Turkey.

SYNONYMS

quadripustulatum Ganglbauer, 1884: 457

Orig. comb.: *Dorcadion quadripustulatum*

Type loc.: Çanakkale (Turkey)

ladikense Breuning & Villiers, 1967: 60

Orig. comb.: *Pedestredorcadion ladikense*

Type loc.: Samsun: Ladik (Turkey)

SPECIES *D. pittinorum* Pesarini & Sabbadini, 1999: 48

Orig. comb.: *Dorcadion pittinorum*

Type loc.: Çorum (Turkey)

Endemic species to Turkey.

- SPECIES** *D. pluto* Thomson, 1867: 47
Orig. comb.: *Dorcadion pluto*
Type loc.: ["Siberia"] clearly mislabeled, should be Turkey
Endemic species to Turkey.
- SPECIES** *D. poleti* Breuning, 1948: 59
Orig. comb.: *Dorcadion poleti*
Type loc.: Caria: Denizli: Babadağ (Turkey)
Endemic species to Turkey.
- SPECIES** *D. praetermissum* Pesarini & Sabbadini, 1999: 47
Orig. comb.: *Dorcadion praetermissum*
Type loc.: Bursa: Uludağ (Turkey)
Endemic species to Turkey.
- SUBSPECIES** *D. p. praetermissum* Pesarini & Sabbadini, 1999: 47
Orig. comb.: *Dorcadion praetermissum praetermissum*
Type loc.: Bursa: Uludağ (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. p. mikhaili* Özdikmen, 2010: 457
Orig. comb.: *Dorcadion praetermissum mikhaili*
Type loc.: Bolu: Abant (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. preissi* Heyden, 1894: 86
Orig. comb.: *Dorcadion preissi*
Type loc.: Kastamonu (Turkey)
Endemic species to Turkey.
- SPECIES** *D. pseudarcivagum* Breuning, 1943: 91
Orig. comb.: *Dorcadion pseudarcivagum*
Type loc.: Amasya (Turkey)
Endemic species to Turkey.
- SPECIES** *D. pseudinfernale* Breuning, 1943: 94
Orig. comb.: *Dorcadion pseudinfernale*
Type loc.: Amanos Mts. (Turkey)
Endemic species to Turkey.
- SPECIES** *D. pseudobithyniense* Breuning, 1962: 465
Orig. comb.: *Dorcadion pseudobithyniense*
Type loc.: Balıkesir (Turkey)
Endemic species to Turkey.
- SPECIES** *D. pseudocinctellum* Breuning, 1943: 90
Orig. comb.: *Dorcadion pseudocinctellum*
Type loc.: Konya: Akşehir (Turkey)
Endemic species to Turkey.
- SPECIES** *D. pseudoholosericum* Breuning, 1962: 300
Orig. comb.: *Dorcadion pseudoholosericum*
Type loc.: Sinop (Turkey)
Endemic species to Turkey.
- SPECIES** *D. pseudopreissi* Breuning, 1962: 411
Orig. comb.: *Dorcadion pseudopreissi*
Type loc.: Sinop (Turkey)
Endemic species to Turkey.
- SPECIES** *D. punctipenne* Küster, 1852: 94
Orig. comb.: *Dorcadion punctipenne*
Type loc.: Anatolia (Turkey)
Endemic species to Turkey.

SYNONYM*distinguendum* Pic, 1931: 10**Orig. comb.:** *Dorcadion olympicum* var. *distinguendum***Type loc.:** İstanbul (Turkey)**SPECIES** *D. punctilicolle* Breuning, 1944: 12**Orig. comb.:** *Dorcadion punctilicolle***Type loc.:** Diyarbakır (Turkey)**Endemic species to Turkey.****SPECIES** *D. pygmaeum* Breuning, 1947: 168**Orig. comb.:** *Dorcadion pygmaeum***Type loc.:** Kurdistan (Turkey)**Endemic species to Turkey.****SPECIES** *D. quadripustulatum* Kraatz, 1873: 88**Orig. comb.:** *Dorcadion quadripustulatum***Type loc.:** Çanakkale (Turkey)**Endemic species to Turkey.****SPECIES** *D. regulare* Pic, 1931: 6**Orig. comb.:** *Dorcadion gallipolitanum* var. *regulare***Type loc.:** Edirne: Keşan (Turkey)**SPECIES** *D. rigattii* Breuning, 1966: 145**Orig. comb.:** *Dorcadion rigattii***Type loc.:** Sivas: Zara (Turkey)**Endemic species to Turkey.****SPECIES** *D. rizeanum* (Breuning & Villiers, 1967: 60)**Orig. comb.:** *Pedestredorcadion rizeanum***Type loc.:** Rize: Sivrikaya (Turkey)**Endemic species to Turkey.****SYNONYM***ikizderense* Breuning, 1973: 54**Orig. comb.:** *Dorcadion ikizderense***Type loc.:** Rize: İkizdere (Turkey)**SPECIES** *D. robustum* Ganglbauer, 1884: 500**Orig. comb.:** *Dorcadion robustum***Type loc.:** Amasya (Turkey)**Endemic species to Turkey.****SPECIES** *D. rolandmenradi* Peks, 1992: 197**Orig. comb.:** *Dorcadion rolandmenradi***Type loc.:** Bursa: Ballısaray, Kütahya (Turkey)**Endemic species to Turkey.****SPECIES** *D. rosti* Pic, 1900: 82**Orig. comb.:** *Dorcadion rosti***Type loc.:** Turkey**SPECIES** *D. rufipenne* Breuning, 1946: 118**Orig. comb.:** *Dorcadion rufipenne***Type loc.:** Kastamonu: Ilgaz Mt. (Turkey)**Endemic species to Turkey.****SUBSPECIES** *D. r. rufipenne* Breuning, 1946: 118**Orig. comb.:** *Dorcadion rufipenne***Type loc.:** Kastamonu: Ilgaz Mt. (Turkey)**Endemic species to Turkey.****SUBSPECIES** *D. r. major* Breuning, 1962: 371**Orig. comb.:** *Dorcadion rufipenne major***Type loc.:** Sinop (Turkey)**Endemic subspecies to Turkey.**

SPECIES *D. rufoapicale* Breuning, 1943: 91 [HN]

Orig. comb.: *Dorcadion rufoapicale*

Type loc.: Bayburt (Turkey)

Endemic species to Turkey.

SYNONYM

rufoapicipenne Breuning, 1946: 125 [RN]

Orig. comb.: *Dorcadion rufoapicipenne*

Type loc.: Bayburt (Turkey)

SPECIES *D. saulcyi* Thomson, 1865: 549

Orig. comb.: *Dorcadion rufoapicale*

Type loc.: "Syria" (mistaken information, probably Turkey)

SUBSPECIES *D. s. saulcyi* Thomson, 1865: 549

Orig. comb.: *Dorcadion rufoapicale*

Type loc.: "Syria" (mistaken information, probably Turkey)

SYNONYM

adanense Pic, 1904: 4

Orig. comb.: *Dorcadion destinoi* var. *adanense*

Type loc.: Adana (Turkey)

subelongatum Pic, 1931: 1

Orig. comb.: *Dorcadion destinoi* var. *subelongatum*

Type loc.: "Akbès" (Turkey: Hatay: Akbez not Syria)

SUBSPECIES *D. s. javeti* Kraatz, 1873: 91

Orig. comb.: *Dorcadion saulcyi javeti*

Type loc.: "Akbès" (Turkey: Hatay: Akbez not Syria)

SYNONYM

destinoi Fairmaire, 1884: 173

Orig. comb.: *Dorcadion destinoi*

Type loc.: "Akbès" (Turkey: Hatay: Akbez not Syria)

tabense Pic, 1931: 1

Orig. comb.: *Dorcadion aleppense* var. *tabense*

Type loc.: Gaziantep (Turkey not Syria)

SPECIES *D. scabricolle* (Dalman, 1817: 174)

Orig. comb.: *Lamia scabricolle*

Type loc.: "Georgia" (mistaken information)

SUBSPECIES *D. s. balikesirensis* Breuning, 1962: 460

Orig. comb.: *Dorcadion scabricolle balikesirensis*

Type loc.: Balıkesir (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. s. caramanicum* Daniel & Daniel, 1903: 332

Orig. comb.: *Dorcadion scabricolle* var. *caramanicum*

Type loc.: Cilician Taurus (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. s. paphlagonicum* Breuning, 1962: 459

Orig. comb.: *Dorcadion scabricolle paphlagonicum*

Type loc.: Kastamonu (Turkey)

Endemic subspecies to Turkey.

SUBSPECIES *D. s. uludaghicum* Breuning, 1970: 98

Orig. comb.: *Dorcadion scabricolle uludaghicum*

Type loc.: Bursa: Uludağ (Turkey)

Endemic subspecies to Turkey.

- SPECIES** *D. schultzei* Heyden, 1894: 86
Orig. comb.: *Dorcadion schultzei*
Type loc.: Mardin (Turkey)
Endemic species to Turkey.
- SPECIES** *D. scrobicolle* Kraatz, 1873: 97
Orig. comb.: *Dorcadion scrobicolle*
Type loc.: Anatolia (Turkey)
Endemic species to Turkey.
- SUBSPECIES** *D. s. scrobicolle* Kraatz, 1873: 97
Orig. comb.: *Dorcadion scrobicolle*
Type loc.: Anatolia (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. s. morulum* Holzschuh, 1965: 42
Orig. comb.: *Dorcadion scrobicolle morulum*
Type loc.: Tokat (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. semibrunneum* Pic, 1903: 170
Orig. comb.: *Dorcadion mniszewi* var. *semibrunneum*
Type loc.: İzmir: Bozdağ (Turkey)
Endemic species to Turkey.
- SUBSPECIES** *D. s. semibrunneum* Pic, 1903: 170
Orig. comb.: *Dorcadion mniszewi* var. *semibrunneum*
Type loc.: İzmir: Bozdağ (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. s. anamasum* Pic, 1934: 33
Orig. comb.: *Dorcadion mniszewi anamasum*
Type loc.: Pisidian Taurus: Isparta: Anamas Mts. (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. semilineatum* Fairmaire, 1866: 273
Orig. comb.: *Dorcadion semilineatum*
Type loc.: İzmir: Bozdağ (Turkey)
Endemic species to Turkey.
- SPECIES** *D. semivelutinum* Kraatz, 1873: 52
Orig. comb.: *Dorcadion semivelutinum*
Type loc.: Sivas (Turkey)
Endemic species to Turkey.
- SPECIES** *D. septemlineatum* Waltl, 1838: 469
Orig. comb.: *Dorcadion septemlineatum*
Type loc.: Turkey
- SUBSPECIES** *D. s. abanti* Braun, 1976: 54
Orig. comb.: *Dorcadion semivelutinum abanti*
Type loc.: Bolu: Abant (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. s. demirciense* Breuning, 1966: 19
Orig. comb.: *Dorcadion septemlineatum demirciense*
Type loc.: Anatolia: ?Manisa: Demirci (Turkey)
Endemic subspecies to Turkey.
- SUBSPECIES** *D. s. novemlineatum* Kraatz, 1873: 61
Orig. comb.: *Dorcadion septemlineatum* var. *novemlineatum*
Type loc.: Bilecik, Eskişehir (Turkey)
Endemic subspecies to Turkey.

SUBSPECIES *D. s. octolineatum* Kraatz, 1873: 61
Orig. comb.: *Dorcadion septemlineatum* var. *octolineatum*
Type loc.: ?İstanbul: Karaköy, Bursa (Turkey)
Endemic subspecies to Turkey.

SUBSPECIES *D. s. septemlineatum* Walth, 1838: 469
Orig. comb.: *Dorcadion septemlineatum*
Type loc.: Turkey

SPECIES *D. sinopense* Breuning, 1962: 299
Orig. comb.: *Dorcadion sinopense*
Type loc.: Sinop (Turkey)
Endemic species to Turkey.

SPECIES *D. sinuatevittatum* Pic, 1937: 6
Orig. comb.: *Dorcadion sinuatevittatum*
Type loc.: Carian Taurus: Denizli: Babadağ (Turkey)
Endemic species to Turkey.

SPECIES *D. sodale* Hampe, 1852: 313
Orig. comb.: *Dorcadion sodale*
Type loc.: Unknown (clearly Turkey)
Endemic species to Turkey.

SUBSPECIES *D. s. sodale* Hampe, 1852: 313
Orig. comb.: *Dorcadion sodale*
Type loc.: Unknown (clearly Turkey)
Endemic subspecies to Turkey.

SUBSPECIES *D. s. trapesunticum* Breuning, 1946: 124
Orig. comb.: *Dorcadion sodale* m. *trapesunticum*
Type loc.: Trabzon (Turkey)
Endemic subspecies to Turkey.

SUBSPECIES *D. s. blumenthali* Breuning, 1966: 20
Orig. comb.: *Dorcadion sodale blumenthali*
Type loc.: Giresun (Turkey)
Endemic subspecies to Turkey.

SPECIES *D. sonjae* Peks, 1993: 5
Orig. comb.: *Dorcadion sonjae*
Type loc.: Elazığ: Elmapınarı and Çaybağ (Turkey)
Endemic species to Turkey.

SPECIES *D. steineri* Holzschuh, 1977: 129
Orig. comb.: *Dorcadion steineri*
Type loc.: Rize: Çamlıhemşin: Hacıvanak plateau (Turkey)
Endemic species to Turkey.

SPECIES *D. subatritarse* Breuning, 1966: 146
Orig. comb.: *Dorcadion subatritarse*
Type loc.: Çorum: Boğazkale (Turkey)
Endemic species to Turkey.

SPECIES *D. subcinctellum* Breuning, 1962: 275
Orig. comb.: *Dorcadion subcinctellum*
Type loc.: Malatya: Yazihan: Hamidiye village (Turkey)
Endemic species to Turkey.

SPECIES *D. subinterruptum* Pic, 1900: 12
Orig. comb.: *Dorcadion divisum* var. *subinterruptum*
Type loc.: Konya (Turkey)
Endemic species to Turkey.

- SPECIES** *D. subsericatum* Pic, 1901: 12
Orig. comb.: *Dorcadion subsericatum*
Type loc.: "Syria" (mistaken information, clearly Turkey: Kastamonu)
Endemic species to Turkey.
- SUBSPECIES** *D. s. subsericatum* Pic, 1901: 12
Orig. comb.: *Dorcadion subsericatum subsericatum*
Type loc.: "Syria" (mistaken information, clearly Turkey: Kastamonu)
Endemic subspecies to Turkey.
- SYNONYM**
paphlagoniense Breuning, 1944: 12
Orig. comb.: *Dorcadion paphlagoniense*
Type loc.: Kastamonu (Turkey)
- SUBSPECIES** *D. s. vulneratum* Pesarini & Sabbadini, 1999: 51
Orig. comb.: *Dorcadion subsericatum vulneratum*
Type loc.: Kastamonu: Masruf pass (Turkey)
Endemic subspecies to Turkey.
- SPECIES** *D. subvestitum* Daniel, 1900: 140
Orig. comb.: *Dorcadion subvestitum*
Type loc.: İzmir: Bozdağ-Ödemiş (Turkey)
Endemic species to Turkey.
- SPECIES** *D. syriense* Breuning, 1943: 94
Orig. comb.: *Dorcadion syriense*
Type loc.: Amanos Mts. (Turkey)
Endemic species to Turkey.
- SPECIES** *D. theophilei* Pic, 1898: 57
Orig. comb.: *Dorcadion theophilei*
Type loc.: Trabzon (Turkey)
Endemic species to Turkey.
- SPECIES** *D. turcicum* Breuning, 1963: 35
Orig. comb.: *Dorcadion turcicum*
Type loc.: Balıkesir: Erdek: Paşalimanı (Turkey)
Endemic species to Turkey.
- SPECIES** *D. ullrichi* Bernhauer, 1988: 102
Orig. comb.: *Dorcadion ullrichi*
Type loc.: Malatya (Turkey)
Endemic species to Turkey.
- SPECIES** *D. wagneri* Küster, 1846: 87
Orig. comb.: *Dorcadion wagneri*
Type loc.: Ağrı: Ağrı Mt. (Turkey)
- SPECIES** *D. weyersii* Fairmaire, 1866: 271
Orig. comb.: *Dorcadion subsericatum*
Type loc.: "Kisilgye-Aole" (Turkey: ?İzmir or Burdur: Kızılkaya)
Endemic species to Turkey.
- SUBGENUS** *MACULATODORCADION* Breuning, 1943: 525
- SPECIES** *D. janssensi* Breuning, 1966: 21
Orig. comb.: *Dorcadion janssensi*
Type loc.: Rize: Tatos Mts. (Turkey)
Endemic species to Turkey.
- SPECIES** *D. quadrimaculatum* Küster, 1848: 79
Orig. comb.: *Dorcadion quadrimaculatum*
Type loc.: Greece and Turkey

SUBSPECIES *D. q. quadrimaculatum* Küster, 1848: 79
Orig. comb.: *Dorcadion quadrimaculatum quadrimaculatum*
Type loc.: Greece and Turkey

SUBSPECIES *D. q. nodicorne* Tournier, 1872: 287
Orig. comb.: *Dorcadion nodicorne*
Type loc.: "Taurus" (Turkey)

SPECIES *D. triste* Frivaldszky, 1845: 184
Orig. comb.: *Dorcadion triste*
Type loc.: "Smyrna" (Turkey: İzmir)
Endemic species to Turkey.

SUBSPECIES *D. t. triste* Frivaldszky, 1845: 184
Orig. comb.: *Dorcadion triste triste*
Type loc.: "Smyrna" (Turkey: İzmir)
Endemic species to Turkey.

SUBSPECIES *D. t. phrygicum* Peks, 1993: 8
Orig. comb.: *Dorcadion triste phrygicum*
Type loc.: Antalya: Ova SE Korkuteli (Turkey)
Endemic species to Turkey.

SPECIES *D. wolfi* Krätschmer, 1985: 24
Orig. comb.: *Dorcadion wolfi*
Type loc.: Balıkesir: Ören (Turkey)
Endemic species to Turkey.

SUBGENUS *MEGALODORCADION* Pesarini & Sabbadini, 1999: 58

SPECIES *D. angorensis* Ganglbauer, 1897: 57
Orig. comb.: *Dorcadion escherichi*
Type loc.: "Angora" (Turkey: Ankara)
Endemic species to Turkey.

SPECIES *D. escherichi* Ganglbauer, 1897: 54
Orig. comb.: *Dorcadion escherichi*
Type loc.: "Angora" (Turkey: Ankara)
Endemic species to Turkey.

SYNONYM

egregium Ganglbauer, 1897: 56
Orig. comb.: *Dorcadion egregium*
Type loc.: "Angora" (Turkey: Ankara)

SPECIES *D. glabrofasciatum* K. Daniel, 1900: 140
Orig. comb.: *Dorcadion glabrofasciatum*
Type loc.: "Bithynia" (Turkey)
Endemic species to Turkey.

SPECIES *D. ledereri* Thomson, 1865: 548
Orig. comb.: *Dorcadion ledereri*
Type loc.: Turkey
Endemic species to Turkey.

SPECIES *D. parallelum* Küster, 1847: 79
Orig. comb.: *Dorcadion parallelum*
Type loc.: Turkey
Endemic species to Turkey.

SPECIES *D. walteri* Holzschuh, 1991: 55
Orig. comb.: *Dorcadion walteri*
Type loc.: Bolu: Alpu Bey village (Turkey)
Endemic species to Turkey.

GENUS *NEODORCADION* Ganglbauer, 1884: 437

SUBGENUS -

SPECIES *N. exornatoides* Breuning, 1962: 68

Orig. comb.: *Neodorcadion exornatoides*

Type loc.: Bahkesir (Turkey)

Endemic species to Turkey.

SPECIES *N. laqueatum* (Waltl, 1838: 469)

Orig. comb.: *Dorcadion laqueatum*

Type loc.: Turkey

SYNONYMS

interruptum Mulsant & Rey, 1863: 150

Orig. comb.: *Dorcadion interruptum*

Type loc.: "Constantinople" (Turkey: İstanbul)

sparsum Mulsant & Rey, 1863: 152

Orig. comb.: *Dorcadion sparsum*

Type loc.: "Constantinople" (Turkey: İstanbul)

SPECIES *N. orientale* Ganglbauer, 1884: 510

Orig. comb.: *Neodorcadion orientale*

Type loc.: Turkey

Endemic species to Turkey.

SPECIES *N. pelleti* (Mulsant & Rey, 1863: 149)

Orig. comb.: *Dorcadion pelleti*

Type loc.: "Smyrna" (Turkey: İzmir)

SYNONYM

segne Mulsant & Rey, 1863: 155

Orig. comb.: *Dorcadion segne*

Type loc.: "Smyrna" (Turkey: İzmir)

As a result of the present work, a total of 204 species group taxa (138 species and 66 subspecies) of Cerambycidae (Dorcadioninae) [*Dorcadion* (134 sp. and 66 ssp.) and *Neodorcadion* (4 sp.)] are determined.

Among of them, 189 species group taxa (130 sp. and 59 ssp.) are endemic to Turkey now. So, endemism ratio of the longhorned beetles that originally described from whole territories of Turkey, is approximately 93 % [Cerambycidae [Dorcadioninae: *Dorcadion* (128 sp. and 59 ssp.) and *Neodorcadion* (2 sp.)]].

LITERATURE CITED

Löbl I. & Smetana A. (ed.) 2010. Catalogue of Palaearctic Coleoptera, Vol. 6. Chrysomeloidea. Stenstrup: Apollo Books, 924 pp.

Özdikmen, H. 2012a. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part I - Vesperidae and Cerambycidae (Prioninae, Lepturinae, Necydalinae, Aseminae, Saphaninae and Dorcasominae). *Munis Entomology & Zoology*, 7 (1): 592-603.

Özdikmen, H. 2012b. The longhorned beetles that originally described from whole territories of Turkey (Coleoptera: Cerambycoidea) Part II - Cerambycidae (Cerambycinae and Stenopterinae). *Munis Entomology & Zoology*, 7 (2): 699-707.

Özdikmen, H. 2012c. Naked lists of Turkish Cerambycoidea and Chrysomeloidea (Coleoptera). *Munis Entomology & Zoology*, 7 (1): 51-108.