

***DORCADION (MACULATODORCADION) PHRYGICUM* PEKS,
1993 STAT. NOV. (CERAMBYCIDAE)**

Hüseyin Özdikmen* and Gamze Kaya*

* Gazi Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, 06500 Ankara / TÜRKİYE. E-mail: ozdikmen@gazi.edu.tr

[Özdikmen, H. & Kaya, G. 2015. *Dorcadion (Maculatodorcadion) phrygicum* Peks, 1993 stat. nov. (Cerambycidae). Munis Entomology & Zoology, 10 (1): 199-200]

ABSTRACT: With the present work, *Dorcadion (Maculatodorcadion) phrygicum* Peks, 1993 stat. nov. is discussed and proposed as a separate species.

KEY WORDS: *Dorcadion (Maculatodorcadion) phrygicum*, stat. nov., Cerambycidae, Turkey.

***Dorcadion phrygicum* Peks, 1993 stat. nov.**

The Turkish endemic species, *Dorcadion triste*, was originally described by Frivaldszky von Frivald (1845) from İzmir province in W Turkey.

Then in 1993, *Dorcadion phrygicum* was originally described by Peks (1993) from Antalya province Kaş county in SW Turkey as a subspecies of *Dorcadion triste* Frivaldszky von Frivald, 1845. Also, it is endemic to Turkey.

Later, it was given by Löbl & Smetana (2010) as a subspecies of *Dorcadion triste* normally. Namely, the South-Eastern populations of *Dorcadion triste* is named *Dorcadion triste phrygicum*.

However, known distribution patterns of *Dorcadion triste triste* and *Dorcadion triste phrygicum* are shown that the taxa do not be subspecies of the same species. As known the South-Eastern populations, *Dorcadion triste phrygicum*, is also distributed in North-Western Anatolia (see below).

Therefore, we propose that *Dorcadion phrygicum* should be as a separate species necessarily.

***Dorcadion triste* Frivaldszky von Frivald, 1845: 184** (Syntypes ♂♂ & ♀♀, ex collection Imre Frivaldszky, Magyar Természettudományi Múzeum, Budapest) [type locality “Smyrna” (Turkey: İzmir)] **A:** TR

vittipenne Breuning, 1946: 97 (*D. triste* m.) [Turkey: İzmir]

postmedioreductum Breuning, 1946: 97 (*D. triste* m.) [Turkey: İzmir]

Known distribution of *D. triste*: İzmir prov. As the type locality (Frivaldszky von Frivald, 1845); Bursa prov. (Ganglbauer, 1884); İzmir prov., Bursa prov. (Aurivillius, 1921); Bursa prov., Balıkesir prov., İzmir prov. (Breuning, 1962); İzmir prov.: Dikili (Makaron) (Demelt, 1963; Gül-Zümreoğlu, 1972); Balıkesir prov.: Balya (Braun, 1978); İzmir prov.: Dikili (Kratschmer, 1985); İzmir prov., Balıkesir prov.: Susurluk, Bursa prov.: Mustafa Kemal Paşa, Karacabey (Önalp, 1990); Anatolia (Danilevsky in Löbl & Smetana, 2010) (Fig. 1).

***Dorcadion phrygicum* Peks, 1993: 8 stat. nov.** (Holotype ♂, collection Heinz Peks, Schwanfeld) [type locality “Kaş: Ova” (Turkey: Antalya)] **A:** TR

Known distribution of *D. phrygicum*: Antalya prov.: Kaş: Ova as the type locality and Bakacak beli N Saklıkent (Peks, 1993); Antalya (Adalia) (Breuning, 1962); Anatolia as *D. triste phrygicum* (Danilevsky in Löbl & Smetana, 2010); Antalya prov.: Aspendos (personal communication with Chris Bruggeman

(Belgium), 2010); Düzce prov.: Kalcı Konutlar, Kent forest road return, Kent forest (Özdikmen et al., 2012); Çanakkale: Biga: Karabiga, Bursa prov.: Uludağ Üniversitesi (Campus of Görükle) (personal data, 2014) (Fig. 2).

LITERATURE CITED

- Aurivillius, C.** 1921. Coleopterorum Catalogus, pars 73, Cerambycidae: Lamiinae. Berlin. W. Junk & S. Schenking. 704 pp.
- Braun, W.** 1978. Die Dorcadienausbeute der Forschungsreisen von W. Heinz 1963-1977, Faunistische Aufstellung, Beschreibung einer neuen Unterart und Bemerkungen zur Systematik wenig bekannter Arten (Coleoptera, Cerambycidae). Nachrichtenblatt der Bayerischen Entomologen, 27 (6): 101-116.
- Breuning, S.** 1962. Revision der Dorcadionini. Entomologische abhandlungen und berichte aus dem staatliche museum für tierkunde in Dresden, 27: 665.
- Demelt, C. V.** 1963. Beitrag zur Kenntnis der Cerambycidenfauna Kleinasien und 13. Beitrag zur Biologie palaearkt. Cerambyciden, sowie Beschreibung einer neuen Oberea-Art. Entomologische Blätter, 59 (3) : 132-151.
- Frivaldszky von Frivald, I.** 1845. A Királyi Magyar természettudományi társulat évkönyvei I. Annalen der ungarischen naturforschenden Gesellschaft, Pesten, 1: 161-187.
- Ganglbauer, L.** 1884. Bestimmungs-Tabellen der europäischen Coleopteren. VIII. Cerambycidae. Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien, 33: 437-586.
- Gül-Zümreoglu, S.** 1972. Catalogue of Insect and common pests (1928-1969). T. C. Publications of Agriculture Ministry, Bornova, İzmir, 119 pp.
- Krätschmer, O.** 1985. Eine neuer *Maculatodorcadion*. Beitrag zur Kenntnis der *Dorcadion*-Fauna Anatoliens (Coleoptera: Cerambycidae: Lamiinae). Entomologische Zeitschrift, 95: 24-27.
- Löbl, I. & Smetana, A. (ed.)** 2010. Catalogue of Palearctic Coleoptera, Vol. 6. Chrysomeloidea. Stenstrup: Apollo Books, 924 pp.
- Önalp, B.** 1990. Systematic researches on *Dorcadion* Dalman, 1817 species in Turkey (Coleoptera, Cerambycidae: Lamiinae) I. H. Ü. Eğitim Fakültesi Dergisi, 5: 57-102.
- Özdikmen, H., Mercan, N. & Tunç, H.** 2012. Longhorned beetles of Düzce province in Turkey (Coleoptera: Cerambycidae). Munis Entomology & Zoology, 7 (2): 714-731.
- Peks, H.** 1993. Drei neue Taxa der Gattung *Dorcadion* aus Anatolien (Coleoptera, Cerambycidae, Lamiinae). Coleoptera. Schwanfelder Coleopterologische Mitteilungen, 1: 1-11.




Figure 1. The distribution of *Dorcadion triste*.




Figure 2. The distribution of *Dorcadion phrygicum*.