

**CONTRIBUTIONS TO THE KNOWLEDGE OF
TURKISH AUCHENORRHYNCHA (HOMOPTERA,
FULGOROMORPHA AND CICADOMORPHA , EXCL.
CICADELLIDAE) WITH A NEW RECORD, *SETAPIUS
KLAPPERICHIANUS* DLABOLA, 1988**

Emine Demir*

* Gazi University, Faculty of Science & Arts, Department of Biology, 06500 Teknikokullar, Ankara, TURKEY. E-mail: eminedemird@yahoo.com

[Demir, E. 2007. Contributions to the knowledge of Turkish Auchenorrhyncha (Homoptera, Fulgoromorpha and Cicadomorpha , excl. Cicadellidae) with a new record, *Setapius klapperichianus* Dlabola, 1988. Munis Entomology & Zoology 2 (1): 39-58]

ABSTRACT: In this study, 2721 samples of Auchenorrhyncha (excl. Cicadellidae) species collected from Antalya region in south-west Turkey during field studies in the years 1999 and 2001-2003 are examined. 74 species belonging to 13 families are found to be distributed in the region. Among these, 14 species belong to the Family Cixidae, 13 species to Delphacidae, 1 species to Meenoplidae, 1 species to Achilidae, 6 species to Dictyopharidae, 1 species to Tropiduchidae, 6 species to Tettigometridae, 2 species to Caliscelidae, 11 species to Issidae, 1 species to Flatidae, 9 species to Cicadidae, 7 species to Cercopidae and 2 species to Membracidae. Distribution of these species in Turkey and their host plants are given along with their locality records. 47 of these examined species are new records for Antalya and *Setapius klapperichianus* Dlabola, 1988 is a new record for Turkey's fauna.

KEY WORDS: Homoptera, Fulgoromorpha, Cicadomorpha, fauna, new record, Turkey.

Turkey's first faunistic records of Auchenorrhyncha (excl. Cicadellidae) are given by Fahringer (1922), Dlabola (1957, 1971a, 1971b, 1974, 1977, 1979a 1979b, 1980, 1981, 1982, 1983a, 1983b, 1984a, 1984b, 1985, 1986, 1995), Linnauvori (1965), Boulard (1979), Asche (1982), Hoch and Remane (1985), Kalkandelen (1980, 1987, 1988) and Kartal (1978, 1980, 1981, 1983, 1985a, 1985b, 1986, 1987). Lodos and Kalkandelen (1980a, 1980b, 1980c, 1981a, 1981b, 1981c, 1988) compile the faunistic records of the studies until 1988 and state that 302 Auchenorrhyncha (excl. Cicadellidae) species are distributed in Turkey, in the checklist they prepared. Later, the studies of Kalkandelen (1989a, 1989b, 1990, 1993, 1994, 2000), Boulard (1993), Kartal, et. al. (1994), Güçlü (1996), Demir (1998, 2006), Emeljanov (2002) and Gnezdilov (2002) present 17 records of species belonging to Auchenorrhyncha (excl. Cicadellidae). According to the studies until recently, 319 species of Auchenorrhyncha (excl. Cicadellidae) were known to be distributed in Turkey.

The province of Antalya, which is selected as the survey area, is located on the Mediterranean coastline in southwest of Turkey, between $29^{\circ}15' - 32^{\circ}36'$ eastern longitudes and $36^{\circ}06' - 37^{\circ}27'$ northern latitudes. The elevation ranges between 0 and 3070m This province is

important because of its biological diversity with four national parks and three nature protection areas; namely, Olympos Beydağları Coastal National Park, Termessos Güllük Mountain National Park, Köprülü Kanyon National Park, Altınbeşik National Park, Çığlıkara Nature Protection Area, Dibek Nature Protection Area and Alacadağ Nature Protection Area. 40 Auchenorrhyncha (excl. Cicadellidae) species are found to be distributed in the survey area, as a result of compilation of the records in the studies of the authors mentioned above.

Climate of the Study Area:

The region is under the influence of a Mediterranean climate that is an outer tropical climate, in which photoperiodism is daily and seasonal, precipitation occurs mostly in cold or relatively colder seasons, summer is the dry period and summer drought coincides with maximum summer temperature. All the stations except Gazipaşa (796.3 mm), which is the easternmost station, receive precipitation over 1000mm, according to Akman (1995). On the coastline, 60-65% of precipitation is received in winter and 0.5-2% in summer.

Vegetation of the Survey Area:

There are communities dominated by *Pinus brutia* forest, garrigue and maquis along the coastline at approximately up to 1000 m above sea level, although they are damaged by human activity in many places. *Pinus brutia* can sometimes be seen above 1000 meters and on the coast it is mixed with *Pinus pinea*, *Ceratonia siliqua*, *Pistacia lentiscus*, *Euphorbia dendroides*, *Myrtus communis*, *Clematis cirrhosa*, *Rubia tinctoria*, *Teucrium creticum* and *Capparis spinosa* are examples of common characteristic species of the coastline. Examples of common characteristic species of the zone just above the coastline are *Quercus coccifera*, *Pistacia terebinthus* ssp. *palaestina*, *Crataegus aronia* ssp. *aronia*, *Fontanesia phillyreoides*, *Rhamnus oleoides* ssp. *graecus*, *Arbutus andrachne*, *Melica eligulata*, *Eryngium falcatum* and *Rubia tenuifolia*. On the eastern half of the region at elevations of 1500m and higher, there are forests comprising of *Cedrus libani*, *Abies cilicica*, *Pinus nigra* ssp. *pallasiana*, *Quercus infectoria* ssp. *boissieri*, *Quercus libani*, *Juniperus excelsa*, and *Juniperus drupacea*. The western parts of the region are covered with *Cedrus libani* forests at elevations of 1500m and higher. Although cedars here show floristic structure, blackpine forests are extremely poor. *Lonicera nummulariifolia* ssp. *glandulifera*, *Digitalis cariensis* and *Acer hyrcanum* ssp. *sphaerocaryum* are examples of characteristic species in this part Akman (1995).

MATERIALS AND METHODS

Field studies in 143 different localities were carried out during the months of March-October in the years 1999 and 2001-2003. In this

study, 2721 adult Auchenorrhyncha (excl. Cicadellidae) samples were collected by sweeping the plants with a wooden shaft net and the of samples collected from only one species of plant were determined. Homopterans, found in the net after sweeping were collected with an aspirator. The samples in the aspirator were then killed by ethyl acetate in a jar and put in standard envelopes and prepared according to the standard methods to produce museum material. The studied samples are preserved in the collection of the author.

RESULTS

There were found to be 74 species of Auchenorrhyncha (excl. Cicadellidae) in the survey area. These are listed below, together with their distribution in Turkey and the plants found to be their host. Out of these examined species 47 are new records for Antalya and *Setapius klapperichianus* Dlabola, 1988 is a new record for Turkey.

Family: Cixiidae Spinola, 1839

***Cixius (Ceratocixius) pallipes* Fieber, 1876**

Materials: Manavgat, Değirmenli, 04.08.1999 1♀ on *Salix*. **Distribution in Turkey:** Adana, Adiyaman, Afyon, Ankara, Antalya, Artvin, Aydın, Balıkesir, Bartın, Çanakkale, Çankırı, Diyarbakır, Düzce Erzincan, Erzurum, Eskişehir, Gaziantep, Giresun, Gümüşhane, Hakkari, İğdır, İzmir, Karaman, Kırşehir, Konya, Kütahya, Kahramanmaraş, Malatya, Mardin, Muğla, Ordu, Sakarya, Samsun, Tokat, Şanlıurfa, Yozgat (Lodos & Kalkandelen, 1980a; Dlabola, 1981; Kalkandelen, 1988). **Remarks:** Known to Antalya.

***Cixius (Ceratocixius) remotus* Edwards, 1888**

Materials: Manavgat, Yukarışıkışıklar-Tilkiler, 460 m., 14.05.2001 1♂1♀ on *Cistus creticus*. **Distribution in Turkey:** Adana, Erzurum, Konya (Kalkandelen, 1988). **Remarks:** New for Antalya.

***Duilius seticulosus* (Lethierry, 1874)**

Materials: Serik: Aspendos, 18.07.2002 14♂2♀; Manavgat: Taşağıl, 60 m., 27.08.2003 11♂5♀; Haciobası, 26.08.2003 6♂ on *Tamarix*. **Distribution in Turkey:** Antalya, Erzincan, Kahramanmaraş, Nevşehir (Lodos & Kalkandelen, 1988; Kalkandelen, 1989). **Remarks:** Known to Antalya.

***Eumecurus gyaurus* (Dlabola, 1957)**

Materials: İbradı: Üzümdere, 470 m., 18.08.1999 1♀; Düzlerçamı, 280 m., 22.07.2001 1♀ on maqius shrubs. **Distribution in Turkey:** Adana, Antalya (Dlabola, 1957; Kalkandelen, 1989b). **Remarks:** Endemic to Turkey and known to Antalya.

***Pentastiridius (s. str.) leporinus* (Linnaeus 1761)**

Materials: Finike, 06.06.2001 1♀. It has been collected from weeds in *Vitex*-shrubs. **Distribution in Turkey:** Antalya, Konya, Mersin, Toros Dağları. (Linnavuori, 1965; Dlabola, 1981; Kalkandelen, 1990). **Remarks:** Known to Antalya.

***Reptalus (s. str.) horridus* (Linnauvori, 1962)**

Materials: Manavgat: Demirciler, 10 m., 17.08.1999 1♂; Demirciler, 40 m., 24.07.2001 1♂1♀; Akseli: Murtıcı, 540 m., 25.07.2001 1♂; İbradı: Başlar, 1440 m., 21.08.2002 2♂2♀ on *Quercus*. **Distribution in Turkey:** Adana, İzmir (Kalkandelen, 1994). **Remarks:** New for Antalya.

***Reptalus oleae* Dlabola, 1987**

Materials: Manavgat: Tilkiler, 470 m., 14.05.2001 1♂; Beşkonak, Köprülü Kanyon, 495 m., 07.06.2001 1♂3♀; Demirciler, 105 m., 18.05.2002 4♂2♀; Demirciler, 105 m., 21.06.2002 1♂1♀; Korkuteli: Korkuteli-Kargalık, 1060 m., 06.06.2001 1♂; Gündoğmuş: Güneycik, 210 m., 22.06.2002 1♀; Akseli: Güçlüköy, 575 m., 22.06.2002 1♀ on *Arbutus andrachne*, *Pinus brutia*, *Celtis australis* and *Verbascum*. **Distribution in Turkey:** Antalya; Kaş, Mersin, Muğla. (Kalkandelen, 1994). **Remarks:** Endemic to Turkey and known to Antalya.

***Setapius klapperichianus* Dlabola, 1988**

Materials: Manavgat: Bucakşeyhler, 24.07.1999 2♀; Manavgat, 21.07.1999 1♀; Manavgat, 10.08.1999 1♂1♀; Manavgat, 10 m., 20.07.1999 1♀; Değirmenli, 04.08.1999 1♀; Demirciler, 65 m., 21.07.1999 1♂; Manavgat, 18.09.1998 2♂3♀; Beşkonak, 145 m., 07.06.2001 2♂; Örenşehir, 15.10.2001 1♀; Çaklış, 16.10.2001 1♂; Manavgat, 10 m., 07.09.2001 1♂4♀; Kemer: Boğürtlenözü, 18.04.2001 1♂3♀; Kemer, 06.06.2001 1♂; Alanya: Okurcalar, 20.05.2002 1♀; Serik: Dikmen, 18.07.2002 1♂; Finike: Yeşilyurt, 18.04.2001 1♂2♀. It has been collected from weeds in fields and *Pinus brutia* forest. **Distribution in Turkey:** This species is the first record in Turkey. **Remarks:** New for Turkey.

***Pentastira major* Kirschbaum, 1868**

Materials: Manavgat: Demirciler, 10 m., 17.08.1999 1♂ on *Quercus*. **Distribution in Turkey:** Adana, Ankara, Antalya; Alanya, Bolu, Çankırı, Çorum, Diyarbakır, Gaziantep, Giresun, Hatay, Isparta, İzmir, Konya, Karaman, Kahramanmaraş, Manisa, Mardin, Muş, Niğde, Samsun, Tekirdağ, Tokat, Şanlıurfa, Van, Yozgat (Linnauvori, 1965; Lodos & Kalkandelen, 1980a; Kalkandelen, 1993). **Remarks:** Known to Antalya.

***Pentastira megista* Emeljanov, 1978**

Materials: İbradı: Ormana, 1000 m., 21.08.2002 1♂ on maquis shrubs. **Distribution in Turkey:** Adana, Ankara, Çankırı, Diyarbakır, Erzurum, Konya, Karaman, Muş, Niğde, Yozgat (Kalkandelen, 1993). **Remarks:** New for Antalya.

***Hyalesthes luteipes* Fieber, 1876**

Materials: Manavgat: Demirciler, 40 m., 17.07.2002 1♂ on *Fontanesia phillyreoides*. **Distribution in Turkey:** Ankara, Aydin, Çorum, Diyarbakır, İstanbul, Kırşehir, Kahramanmaraş, Karaman, Kayseri, Konya, Mersin, Nevşehir, Van, Yozgat (Linnauvori, 1965; Dlabola, 1957, 1981; Lodos & Kalkandelen, 1980a; Hoch & Remane, 1985; Kalkandelen, 2000). **Remarks:** New for Antalya.

***Hyalesthes mavromoustakisi* Dlabola 1959**

Materials: Gündoğmuş: Serinyaka, 700 m., 25.07.1999, 1♀; Korkuteli: Korkuteli, 30 km W, 670 m., 28.08.2003 2♀ on weeds and maquis shrubs.
Distribution in Turkey: Diyarbakır (Lodos & Kalkandelen, 1980a).
Remarks: New for Antalya.

***Hyalesthes mlokosiewiczi* Signoret, 1879**

Materials: Korkuteli: Koru Dağı, Kartal Yaylası, 1600 m., 23.06.2002 1♀ on *Salix*. **Distribution in Turkey:** Adana, Adiyaman, Afyon, Ankara, Aydin, Bilecik, Burdur, Çankırı, Diyarbakır, Gaziantep, Isparta, İzmir, Kahramanmaraş, Kilis, Malatya, Mardin, Mersin, Muğla, Siirt, Şanlıurfa, Tokat (Dlabola, 1957, 1981; Lodos & Kalkandelen, 1980a, Kalkandelen, 2000).

Remarks: New for Antalya.

***Hyalesthes obsoletus* Signoret, 1865**

Materials: İbradı: Üzümdere, 470 m., 18.08.1999, 3♂2♀; Manavgat: Demirciler 40 m, 11.05.2001 1♀; Haciobası, 14.10.2001 1♂; Haciobası, 22.06.2002 1♀; Taşgil, 60 m., 27.08.2003 4♂; Haciobası, 26.08.2003 4♂; Akseki: Erenkaya, 630 m., 22.06.2002 3♂2♀; Finike: Dağbağ, 475 m., 13.07.2002 1♂1♀; Kale: Kale, 20.08.2002 1♀. It has been collected from weeds in fields and *Vitex agnus-castus*. **Distribution in Turkey:** Adana, Afyon, Adiyaman, Ağrı, Ankara, Antalya, Aydin, Balıkesir, Bolu, Burdur, Bursa, Çanakkale, Çankırı, Çorum, Diyarbakır, Düzce, Edirne, Elazığ, Erzincan, Erzurum, Eskişehir, Gaziantep, Giresun, Hakkari, İğdır, Isparta, , İstanbul, Kahramanmaraş, Konya, Malatya, Mardin, Mersin, Muğla, Nevşehir, Ordu, Rize, Sakarya, Sinop, Sivas, Tokat, Trabzon, Şanlıurfa, Van (Fahringer, 1922; Dlabola, 1957, 1981; Hoch & Remane, 1985; Lodos & Kalkandelen, 1980a; Kalkandelen, 2000). **Remarks:** Known to Antalya.

Family: Delphacidae Leach, 1815***Asiraca clavicornis* (Fabricius, 1794)**

Materials: Manavgat: Evrenleryavşı, 10 m., 16.03.2002 1♀. It has been collected from weeds in *Pinus brutia*. **Distribution in Turkey:** Amasya, Ankara, Antalya, Aydin, Çorum, Denizli, Erzurum, İstanbul, İzmir, Konya, Kütahya, Muğla, Nevşehir, Samsun, Sinop, Yozgat (Fahringer, 1922; Lodos & Kalkandelen, 1980b; Dlabola, 1981; Asche, 1982; Güçlü, 1996). **Remarks:** Known to Antalya.

***Kelisia ribauti* Wagner, 1938**

Materials: Akseki: Geriş, 700 m., 17.04.2001 7♂11♀. It has been collected from weeds in *Pinus brutia*. **Distribution in Turkey:** Ankara, Aydin, Bitlis, Denizli, Diyarbakır, Erzurum, Hakkari, Konya, Malatya, Siirt, Van (Dlabola, 1957, 1981; Lodos & Kalkandelen, 1980b; Asche, 1982; Güçlü, 1996). **Remarks:** New for Antalya.

***Kelisia yarkonensis* Linnauvori, 1962**

Materials: Manavgat: Boztepe, Alara, 15.10.2001 1♀; Bucakşeyhler, 70 m., 23.04.2002 1♂; Kale, Eseler, 55 m, 15.05.2001 1♀. It has been collected from weeds in fields. **Distribution in Turkey:** Konya (Asche, 1982). **Remarks:** New for Antalya.

***Tropidocephala tuberipennis* (Mulsant et Rey, 1855)**

Materials: Manavgat: Bucakşeyhler, 24.07.1999 1♀; Bucakşeyhler, 13.08.1999 2♀; Haciobası, 14.10.2001 1♂; Demirciler, 40 m., 20.06.2002 43♂35♀; Kemer: Kemer, 06.09.2002 1♀. On Graminae. **Distribution in Turkey:** Adana, Antalya, Aydin, Mersin (Dlabola, 1957; Asche, 1982). **Remarks:** Known to Antalya.

***Chloriana ponticana* Asche, 1982**

Materials: Gazipaşa: Çalıpinar-Kahyalar, 20 m., 20.05.2002 1♀ on *Arundo donax*. **Distribution in Turkey:** Antalya (Asche, 1982). **Remarks:** Known to Antalya.

***Chloriana flaveola* Lindberg, 1948**

Materials: Manavgat: Demirciler, 65 m., 15.04.2001 1♀; Kumluca: Kumluca, 06.06.2001 6♀; Alanya: Konaklı, 25 m, 24.04.2002 1♀; Düzlerçami: 280 m., 22.07.2001 1♀ on *Arundo donax*. **Distribution in Turkey:** Adana (Dlabola, 1957). **Remarks:** New for Antalya.

***Laodelphax striatellus* (Fallen, 1826)**

Materials: Manavgat: Bucakşeyhler, 13.08.1999 1♂2♀; Bucakşeyhler, 18-28.06.1997 1♀; Beşkonak, 145 m., 07.06.2001 5♂2♀; Demirciler, 65 m., 05.06.2001 1♂; Demirciler, 20 m., 18.05.2002 2♂2♀; Demrciler, 40 m, 18.08.2001 6♂13♀; Manavgat, 10 m, 07.09.2001 3♂5♀; Alanya: Obaköy, 25 m., 12.08.1999 1♂3♀; Elmali: Akçay, 1100 m., 28.07.1999 1♂; Kemer: Kemer, 06.06.2001 8♂4♀; Bögürtlenözü, 18.04.2001 5♂8♀; Çamyuva, 18.04.2001 2♂11♀; Kemer, 06.09.2002 1♂1♀; Finike: Turunçova, 50 m., 06.06.2001 6♂13♀; Serik: Aspendos, Köprüçaylı, 16.10.2001 2♀; Camili, 26.07.2001 6♂4♀; Güllük Dağı: Araştırma Ormanı, 750 m., 21.05.2002 1♂; Korkuteli: Datköy, 960 m., 13.07.2002 15♂16♀; Kale: Köşkerler, 240 m, 15.05.2001 3♂9♀; Kumluca, Kavaklıbibi, 130 m, 24.06.2002 7♂; Gazipaşa: Macarköy, 09.09.2001 5♂1♀; Kaş: Kalkan, Üzümlü, 285 m, 15.05.2001 4♂5♀. It has been collected from weeds in *Pinus brutia* forest and fields. **Distribution in Turkey:** Adana, Adiyaman, Ankara, Antalya, Bilecik, Bitlis, Diyarbakır, Erzincan, Erzurum, İğdır, İzmir, Kahramanmaraş, Kars, Malatya, Malatya, Mersin, Muğla, Nevşehir, Niğde, Ordu, Rize, Siirt (Dlabola, 1971, 1981; Lodos & Kalkandelen, 1980b; Asche, 1982; Güçlü, 1996). **Remarks:** Known to Antalya.

***Sogatella vibix* (Haupt, 1922)**

Materials: Kaş: Kalkan, Yeşilköy, 29.07.1999 1♂; Manavgat: Bucakşeyhler, 15-26.08.1997 1♂; Çaklış, 16.10.2001 2♀; Örenşehir, 15.10.2001 1♀(Graminae); Taşağıl, Kılımli, 40 m, 08.09.2002 1♂5♀; İlica, 07.09.2001 2♀; Demirciler, 40 m, 18.08.2001 2♂; Gazipaşa: Çalıpinar-Kahyalar, 20 m., 20.05.2002 1♀; Asarcık, 24.04.2002 3♂18♀; Finike: Yeşilyurt, 18.04.2001 3♂1♀; Erenkavak, 65 m, 24.06.2002 1♂4♀; Yeşilyurt, 06.09.2002 3♂7♀; Serik: Dikmen, 26.07.2001 1♂8♀; Aksu: Topallar, 26.07.2001 2♂13♀; Kale: Eseler, 55 m, 24.06.2002; 6♂15♀; Kemer: Bögürtlenözü, 21.08.2001 5♂2♀; Kemer, 06.09.2002 2♂; Alanya: Konaklı, 25 m, 24.04.2002 4♂. It has been collected from weeds in *Pinus brutia* forest and fields. **Distribution in Turkey:** Ankara, Aydin, Bitlis, Diyarbakır, Erzurum, Hatay, Mersin, Van (Dlabola, 1971, 1981; Asche, 1982; Güçlü, 1996). **Remarks:** New for Antalya.

***Muirodelphax aubei* (Perris, 1857)**

Materials: Elmalı: Çığlıkara, 1600 m., 28.08.2003 3♂3♀. It has been collected from weeds in *Cedrus libani*. **Distribution in Turkey:** Ankara, Denizli, Erzurum, İzmir, Van (Dlabola, 1957, 1981; Lodos & Kalkandelen, 1980b; Asche, 1982; Güçlü, 1996). **Remarks:** New for Antalya.

***Toya propinquua* (Fieber, 1866)**

Materials: Manavgat: 21.07.1999 1♂1♀; Bucakşeyhler, 13.08.1999 2♂1♀; Bucakşeyhler, 24.07.1999 1♂2♀; Manavgat, 18.09.1998 1♂; Demirciler, 10 m., 17.08.1999 1♀; Belenobası, 130 m., 15.04.2001 2♀; Beşkonak, 145 m., 07.06.2001 3♂9♀; Demirciler, 40 m., 05.06.2001 1♂; Bucakşeyhler, 70 m., 23.04.2002 2♂; Demirciler, 20 m., 18.05.2002 1♂; Demirciler, 105 m., 21.06.2002 1♀; Demirciler, 40 m., 20.06.2002 3♀; Haciobası, 22.06.2002 1♂1♀; Alanya: Obaköy, 25 m., 12.08.1999 1♂3♀; Okurcalar, 12.05.2001 1♂; Okurcalar, 09.09.2001 2♂2♀; Konaklı, 25 m, 24.04.2002 3♂2♀; Kaş: Kalkan, Yeşilköy, 29.07.1999 3♂3♀; Finike: Turunçova, 50 m., 06.06.2001 1♂; Kumluca: Kavakköy, 18.04.2001 2♂6♀; Kale: Eseler, 55 m, 15.05.2001 1♂3♀; Serik: Dikmen, 26.07.2001 1♂7♀; Gazipaşa. Asarcık, 24.04.2002 4♂9♀; Kemer: Karabucak, 85 m, 25.05.2002 1♂4♀; Kemer, 06.09.2002 1♂8♀. It has been collected from weeds in *Pinus brutia* and fields. **Distribution in Turkey:** Adana, Afyon, Amasya, Ankara, Antalya, Aydın, Çanakkale, Denizli, Diyarbakır, Erzurum, Gaziantep, Hatay, Kastamonu, Mardin, Mersin, Muğla, Ordu, Samsun, Siirt, Sinop (Dlabola, 1957, 1971; Linnauvori, 1965; Lodos & Kalkandelen, 1980b; Asche, 1982; Güçlü, 1996). **Remarks:** Known to Antalya.

***Pseudaraeopus lethierryi* (Mulsant et Rey, 1879)**

Materials: Manavgat. Bucakşeyhler, 24.07.1999 1♂; Örenşehir, 15.10.2001 1♀; Düzlerçami: 280 m., 22.07.2001 1♀; Gazipaşa: Çalıpinar-Kahyalar, 20 m., 20.05.2002 1♀; Macarköy, 09.09.2001 1♀; Kemer: Olimpos Milli Parkı, 20.08.2002 1♀; Finike, Erenkavak, 65 m, 24.06.2002 1♀. It has been collected from weeds in fields. **Distribution in Turkey:** Aydın (Asche, 1982). **Remarks:** New for Antalya.

***Delphacodoides anaxarchi* (Muir, 1926)**

Materials: Manavgat, Bucakşeyhler, 13.08.1999 4♀; Alanya, Kargıcak, 09.09.2001 3♀; Serik, Camili, 20.08.2001 1♂1♀; Manavgat, Demirciler, 40 m, 18.08.2001 1♀. It has been collected from weeds in *Pinus brutia* forest. **Distribution in Turkey:** Antalya, Mersin (Dlabola, 1957, 1981; Asche, 1982).

Remarks: Known to Antalya.

***Perkinsiella dorsata* (Melichar, 1905)**

Materials: Manavgat, 18.09.1998 1♂; Manavgat, 10.08.1999 1♂; Demirciler, 40 m., 20.06.2002 2♀; Karaöz, 70 m, 18.08.2001 1♂; Kemer, Çamyuva, 18.04.2001 3♀; Aksu, Topallar, 26.07.2001 1♂2♀; Alanya, Okurcalar, 09.09.2001 2♀. It has been collected from weeds in *Pinus brutia* forest. **Distribution in Turkey:** Mersin, Muğla (Kalkandelen, 1980; Dlabola, 1981; Asche, 1982). **Remarks:** New for Antalya.

Family: Meenoplidae Fieber, 1872***Meenoplus albosignatus* Fieber, 1866**

Materials: Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 1♂; Gündoğmuş, Serinyaka, 625 m., 22.06.2002 1♂; Gazipaşa, Macarköy,

22.08.2002 1♀; İbradı, Başlar, 1440 m., 21.08.2002 1♂3♀ on *Quercus*.
Distribution in Turkey: Adiyaman, Ankara, Bolu, Gaziantep, Hakkari, Malatya, Mardin, Muş (Dlabola, 1957, 1981; Linnavuori, 1965; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

Family: Achilidae Stal, 1866

***Cixidia pilatoi* D'Urso & Guglielmino, 1995**

Materials: Manavgat, Demirciler, 105 m., 18.05.2002 2♂3♀ on *Pinus brutia* and *Myrthus communis*. **Distribution in Turkey:** Adana (Linnavuori, 1965). **Remarks:** New for Antalya. The record of Adana has been reported as *C. marginicollis*. This record possibly belongs to *C. pilatoi*.

Family: Dictyopharidae Spinola, 1839

***Callodictya krueperi* (Fieber, 1876)**

Materials: Gündoğmuş, Çiçekoluk, 960 m., 25.07.1999 1♀; Akseki, Murtiçi, 540 m., 25.07.2001 5♂7♀; Manavgat, Demirciler, 40 m., 24.07.2001 2♂; Demirciler, 40 m., 17.07.2002 1♂; Serik, Sarıabahı, 18.07.2002 1♀; Kaş, Kemerköy, 260 m., 13.07.2002 2♀; Dirgenler, 265 m., 13.07.2002 1♂; İbradı, Başlar, 1440 m., 21.08.2002 1♂3♀ on *Quercus*, *Myrthus communis* and *Centaurea*. **Distribution in Turkey:** Hatay, İzmir, Tunceli (Linnavuori, 1965; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Dictyophara (s. str.) asiatica* Melichar, 1912**

Materials: Kaş, Beldibi, 1400 m., 29.07.1999 1♂3♀; Manavgat, Bucakşeyhler, 45 m., 24.07.1999 1♂; Demirciler, 10 m., 17.08.1999 1♂1♀; Bucakşeyhler, 13.08.1999 1♀; Alanya, Obaköy, 25 m., 12.08.1999 1♀; Gündoğmuş, Çiçekoluk, 960 m., 25.07.2001 8♂17♀; Akseki, Murtiçi, 540 m., 25.07.2001 4♀; Manavgat, Demirciler, 40 m., 24.07.2001 3♂5♀; Bucakşeyhler, 45 m., 19.07.2001 3♂1♀; Alanya, Kargıcak, 21.07.2001 2♂; Manavgat, Demirciler, 20 m., 21.06.2002 1♀; Demirciler, 40 m., 20.06.2002 1♂2♀; Demirciler, 40 m., 17.07.2002 1♀; Kaş, Dirgenler, 265 m., 13.07.2002 1♀; Güllük Dağı, Araştırma Ormanı, 750 m., 19.07.2002 1♂; Serik, Dikmen, 18.07.2002 1♀; Hisarçandır, 770 m., 20.08.2002 1♂1♀; Kemer, Çamyuva, 20.08.2002 1♀; Olimpos Milli Parkı, 20.08.2002 1♂1♀; Sarıçınar Dağı, 20.08.2002 2♀; Manavgat, Haciobası, 26.08.2003 1♀ on *Verbascum*, *Pistacia*, *Centaurea*, *Phlomis*, *Fontanesia phillyreoides*, *Quercus*, *Ficus carica*, *Myrthus communis*, *Vitex agnus-castus* and *Populus*. **Distribution in Turkey:** Adana, Amasya, Ankara, Aydın, Balıkesir, Bilecik, Bolu, Burdur, Denizli, Diyarbakır, Elazığ, Eskişehir, Erzincan, Isparta, İzmir, Kırklareli, Kütahya, Manisa, Muğla, Sakarya, Tunceli, Urfa, Van (Dlabola, 1957, 1981; Linnavuori, 1965; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Dictyophara iranica* Linnavuori, 1962**

Materials: Manavgat, Boztepe, Alara, 15.10.2001 1♀; Serik, Aspendos, Köprüçayı, 16.10.2001 1♀; Aspendos, 18.07.2002 1♂; Taşağıl, 60 m., 27.08.2003 1♂2♀ on *Tamarix*. **Distribution in Turkey:** Siirt (Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Dictyophara (s. str.) lindbergi* Metcalf, 1955**

Materials: Manavgat, 10 m., 20.07.1999 1♀; Serik, Dikmen, 18.07.2002 2♂1♀; Akseki, Güneykaya, 570 m., 25.07.1999 1♂1♀; Güllük Dağı, Araştırma Ormanı,

550 m., 19.07.2002 3♂1♀. It has been collected from weeds in *Pinus brutia* forest, *Ficus carica* and *Salix*. **Distribution in Turkey:** Adana, İzmir, Kütahya, Muğla, Tekirdağ (Dlabola, 1957; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Dictyophara (Euthremma) multireticulata* Mulsant et Rey, 1855**

Materials: İbradı, Başlar, 1440 m., 21.08.2002 1♀ on *Quercus*. **Distribution in Turkey:** Denizli, Van (Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Dictyophara (Chanithus) xiphias* Puton, 1884**

Materials: Manavgat, Demirciler, 10 m., 17.08.1999 1♂2♀; Demirciler, 40 m., 21.07.1999 1♀; Gündoğmuş, Çiçekoluk, 960 m., 25.07.2001 1♀; Manavgat, Bucakşeyhler, 45 m., 19.07.2001 1♂; Güllük Dağı, Araştırma Ormanı, 750 m., 19.07.2002 1♂3♀; Gazipaşa, Macarköy, 22.08.2002 2♀. On *Phlomis*, *Fontanesia phillyreoides*, *Pistacia terebinthus*, and *Quercus*. **Distribution in Turkey:** İzmir (Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

Family: Tropiduchidae Stal, 1866

***Trypetimorpha occidentalis* Huang et Bourgoin, 1993**

Materials: Düzlerçami, 280 m., 22.07.2001 1♀. It has been collected from weeds in *Pinus brutia* forest. **Distribution in Turkey:** Ankara (Demir, 1998). **Remarks:** New for Antalya.

Family: Tettigometridae Germar, 1821

***Tettigometra (Hystrigonia) hexaspina* Kolenati, 1857**

Materials: Korkuteli, Koru Dağı, Kartal Yaylası, 1600 m., 23.06.2002 1♂1♀; Söğütcük, 1010 m., 13.07.2002 1♀. It has been collected from weeds in fields. **Distribution in Turkey:** Ağrı, Ankara, Edirne, Gaziantep, Giresun, Isparta, Tekirdağ, Urfa (Dlabola, 1957; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Tettigometra (Metroplaca) longicornis* (Signoret, 1866)**

Materials: Manavgat, Yukarıışıklar-Tilkiler, 460 m., 14.05.2001 1♂1♀; Korkuteli, Koru Dağı, Kartal Yaylası, 1600 m., 23.06.2002 1♂4♀. It has been collected from weeds in fields. **Distribution in Turkey:** Ankara, Sivas (Dlabola, 1957, 1981). **Remarks:** New for Antalya.

***Tettigometra (Mitricephalus) eremi* Lindberg, 1948**

Materials: Manavgat, Yukarıışıklar-Tilkiler, 460 m., 14.05.2001 11♂18♀; Güllük Dağı, Araştırma Ormanı, 750 m., 21.05.2002 3♂4♀; Korkuteli, Koru Dağı, Kartal Yaylası, 1600 m., 23.06.2002 8♂14♀. It has been collected from weeds in *Pinus brutia* forest, fields and *Salix*. **Distribution in Turkey:** Ankara, Aydın, Burdur, Çanakkale, İzmir, Kayseri, Kocaeli, Kütahya, Manisa, Sivas, Uşak (Lodos & Kalkandelen, 1980c; Dlabola, 1981). **Remarks:** New for Antalya.

***Tettigometra (Mitricephalus) leucophaea* (Preyssler 1792)**

Materials: Korkuteli, Söğütcük, 1010 m., 13.07.2002 5♂6♀; İbradı, Başlar, 1440 m., 21.08.2002 1♀; Elmalı, Çığlıkara, 1600 m., 28.08.2003 1♀. It has been collected from weeds in *Juniperus-Abies cilicia* forest and *Quercus*.

Distribution in Turkey: Adana, Ankara, Adiyaman, Ağrı, Bilecik, Bolu, Burdur, Çanakkale, Çankırı, Çorum, Diyarbakır, Edirne, Elazığ, İstanbul, İzmir, Mardin, Nevşehir, Sivas, Tekirdağ, Urfa (Dlabola, 1957; Linnavuori, 1965; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Tettigometra (s. str.) sulphurea* Mulsant et Rey, 1855**

Materials: Manavgat, Belenobası, 130 m., 15.04.2001 1♀; Bucakşeyhler, Seleukeia, 250 m., 23.04.2002 1♀; Bucakşeyhler, 70 m., 23.04.2002 1♂1♀. It has been collected from weeds in *Pinus brutia* forest. **Distribution in Turkey:** Adana, Ankara, Artvin, Aydin, Bilecik, Bursa, Diyarbakır, Elazığ, İzmir, Kütahya, Nevşehir, Sakarya, Urfa, Uşak, Van (Dlabola, 1957; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

***Tettigometra (s. str.) virescens* (Panzer, 1799)**

Materials: Kemer, Ovacık, 1150 m., 20.08.2002 1♂. It has been collected from weeds in *Cedrus libani* forest. **Distribution in Turkey:** Adana, Adiyaman, Ankara, Bursa, Diyarbakır, Elazığ, Erzincan, Gaziantep, Hatay, İğdır, Mardin, Tunceli, Urfa, Van (Fahringer, 1922; Lodos & Kalkandelen, 1980c). **Remarks:** New for Antalya.

Family: Caliscelidae Amyot & Serville, 1834

***Bruchoscelis peculiaris* (Horvath, 1904)**

Materials: Manavgat, Demirciler, 40 m., 20.06.2002 1♀. On Graminae. **Distribution in Turkey:** Ankara, Yozgat (Dlabola, 1957; Kartal, 1985). **Remarks:** New for Antalya and endemic to Turkey.

***Ommatidiotus longiceps* Puton, 1896**

Materials: Serik, Dikmen, 18.07.2002 1♀. It has been collected from weeds in *Pinus brutia* forest. **Distribution in Turkey:** Diyarbakır, Mardin, Urfa (Lodos & Kalkandelen, 1988). **Remarks:** New for Antalya.

Family: Issidae Spinola, 1839

***Mycteroodus (s. str.) lodosicus* Dlabola, 1980**

Materials: Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 3♀; Güllük Dağı, 540 m., 22.07.2001 2♂; Araştırma Ormanı, 550 m., 19.07.2002 1♂; Araştırma Ormanı, 750 m., 19.07.2002 1♂ on *Phlomis*, *Quercus* and *Cytisus laburnum*. **Distribution in Turkey:** Aydin (Dlabola, 1980). **Remarks:** New for Antalya and endemic to Turkey.

***Mycteroodus (Comporodus) spinicordatus* (Dlabola, 1983)**

Materials: Manavgat, Beşkonak, Köprülü Kanyon, 495 m., 07.06.2001 10♂6♀; Gündoğmuş, Güneycik, 210 m., 08.06.2001 1♂; Gazipaşa, Kahyalar, 22.08.2002 1♂; Manavgat, Beşkonak, Köprülü Kanyon, 495 m., 27.08.2003 1♂ on *Arbutus andrachne*, *Ceratonia siliqua* and *Nerium oleander*. **Distribution in Turkey:** Antalya (Dlabola, 1983). **Remarks:** Endemic to Turkey and known to Antalya.

***Mycteroodus (Comporodus) tekneticus* (Dlabola, 1982)**

Materials: Manavgat, Yukarıışıklar-Tilkiler, 460 m., 14.05.2001 3♂1♀; Akseki, Güclüköy, 550 m., 21.04.2002 2♂; Gündoğmuş, Güneycik, 210 m., 21.04.2002 7♂1♀; Güneycik, 240 m., 21.04.2002 2♂1♀; Çiçekoluk, 960 m., 19.05.2002

1♂3♀; Güneycik, 210 m., 22.06.2002 3♂; Akseki, Erenkaya, 630 m., 22.06.2002 1♀; Kemer, Ovacık, 1150 m., 20.08.2002 1♀ on *Phlomis*, *Fontanesia phillyreoides*, *Cercis*, *Arbutus andrachne* and *Quercus*. **Distribution in Turkey:** Antalya (Dlabola, 1982). **Remarks:** Endemic to Turkey and known to Antalya.

***Tshurtshurnella alanyana* Dlabola, 1982**

Materials: Gündoğmuş, Çiçekoluk, 960 m., 25.07.1999 8♂17♀; Manavgat, Bucakşeyhler, 24.07.1999 1♂; Demirciler, 40 m., 21.07.1999 2♂; Demirciler, 40 m., 05.06.2001 1♂; Taşağıl, Kılımli, 40 m., 03.06.2001 2♂3♀; Gündoğmuş, Çiçekoluk, 960 m., 25.07.2001 17♂22♀; Alanya, Keşefli, 21.07.2001 4♂2♀; Manavgat, Bucakşeyhler, 45 m., 19.07.2001 1♀; Demirciler, 40 m., 24.07.2001 1♂; Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 1♂; Manavgat, Demirciler, 40 m., 20.06.2002 6♂7♀; Akseki, Erenkaya, 630 m., 22.06.2002 1♂1♀; Gündoğmuş, Güneycik, 210 m., 22.06.2002 1♂; Güneycik, 210 m., 14.07.2002 1♀; Güllük Dağı, Araştırma Ormanı, 750 m., 19.07.2002 2♀; İbradı, Başlar, 1440 m., 21.08.2002 1♂1♀; Hisarçandır, 770 m., 20.08.2002 1♂3♀; Kemer, Sarıçınar Dağı, 1245 m., 20.08.2002 12♂7♀ on *Verbascum*, *Centaurea*, *Phlomis*, *Capparis*, *Fontanesia phillyreoides*, *Quercus*, *Graminae*, *Cistus creticus* and *Euphorbia*. **Distribution in Turkey:** Antalya (Dlabola, 1982). **Remarks:** Endemic to Turkey and known to Antalya.

***Latematium cingulatum* Dlabola, 1983**

Materials: Manavgat, Beşkonak, Köprülü Kanyon, 495 m., 07.06.2001 1♀; Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 10♂4♀; Güllük Dağı, 540 m., 22.07.2001 2♂; Termessos, Güllük Dağı, 900-1000 m., 21.05.2002 1♂1♀; Güllük dağı, 900-1000 m., 19.07.2002 5♂3♀; Araştırma Ormanı, 550 m., 19.07.2002 1♂; Çığlık, 330 m., 19.07.2002 1♂ on *Arbutus andrachne*, *Cytisus laburnum* and shrubs. **Distribution in Turkey:** Antalya, Kütahya (Dlabola, 1983). **Remarks:** Endemic to Turkey and known to Antalya.

***Latilica antalyica* (Dlabola, 1986)**

Materials: Kaş, Kemerköy, 260 m., 13.07.2002 2♂2♀ on *Arbutus andrachne*. **Distribution in Turkey:** Antalya, Hatay (Dlabola, 1986). **Remarks:** Endemic to Turkey and known to Antalya.

***Latilica maculipes* (Melichar, 1906)**

Materials: Manavgat, Demirciler, 40 m., 24.07.2001 1♂1♀; Gündoğmuş, Çiçekoluk, 960 m., 25.07.2001 1♀; Serik, Sarıabali, 18.07.2002 2♂1♀; İbradı, Başlar, 1440 m., 21.08.2002 1♀ on *Quercus*, *Myrthus communis* and maqius shrubs. **Distribution in Turkey:** Antalya, Hatay, Manisa, Muğla (Linnauvori, 1965; Dlabola, 1971, 1986; Lodos & Kalkandelen, 1981a). **Remarks:** Known to Antalya and known to Antalya.

***Latilica oertzeni* Matsumura, 1910**

Materials: Hisarçandır, 770 m., 20.08.2002 1♀; Kemer, Çamyuva, 20.08.2002 1♀. On *Cistus creticus* ve *Myrthus communis*. **Distribution in Turkey:** Antalya (Lodos & Kalkandelen, 1988). **Remarks:** Known to Antalya.

***Latilica retamae* (Linnauvori, 1962)**

Materials: Manavgat, Demirciler, 65 m., 14.04.2001 1♀; Demirciler, 40 m., 24.07.2001 1♂; Akseki, Murtıçı, 540 m., 25.07.2001 1♂ on *Quercus* and maqius

shrubs. **Distribution in Turkey:** Antalya (Lodos & Kalkandelen, 1988).
Remarks: Known to Antalya.

***Agalmatium bilobum* (Fieber, 1877)**

Materials: Korkuteli, Yazır-İmrähör, 950 m., 27.07.1999 1♀; Akseki, Erenkaya, 680 m., 25.07.1999 1♀; Alanya, Okurcalar, 12.05.2001 1♂1♀; Gazipaşa, Çalıpinar, 25 m., 12.05.2001 6♂6♀; Manavgat, Tilkiler, 470 m., 14.05.2001 1♂1♀; Yukarıışıklar, 200 m., 14.05.2001 16♂15♀; Demirciler, 105 m., 11.05.2001 39♂2♀; Taşağıl, Kilimli, 40 m., 03.06.2001 1♂2♀; Beşkonak, 145 m., 07.06.2001 5♂2♀; Taşağıl, Kilimli, 60 m., 03.06.2001 7♂5♀; Taşağıl, Sağırin, 40 m., 03.06.2001 3♂; Taşağıl, 55 m., 07.06.2001 1♂; Demirciler, 40 m., 05.06.2001 2♂2♀; Demirciler, 65 m., 05.06.2001 1♂1♀; Kemer, 06.06.2001 1♀; Gündoğmuş, Çiçekoluk, 960 m., 25.07.2001 1♂3♀; Akseki, Murtiçi, 540 m., 25.07.2001 2♂; Manavgat, Demirciler, 105 m., 22.04.2002 1♂; Güllük Dağı, Araştırma Ormanı, 750 m., 21.05.2002 6♂; Alanya, Okurcalar, 20.05.2002 1♂; Gazipaşa, Çalıpinar-Kahyalar, 20 m., 20.05.2002 11♂18♀; Manavgat, Demirciler, 105 m., 18.05.2002 20♂40♀; Demirciler, 20 m., 21.06.2002 2♂; Demirciler, 105 m., 21.06.2002 2♂3♀; Demirciler, 40 m., 20.06.2002 2♀; Gündoğmuş, Güneycik, 210 m., 22.06.2002 5♂5♀; Akseki, Erenkaya, 630 m., 22.06.2002 10♂11♀; Elmalı, Gökpınar, 1200 m., 13.07.2002 1♂4♀; Akçay, 1210 m., 13.07.2002 2♂; Güllük Dağı, Araştırma Ormanı, 550 m., 19.07.2002 2♂1♀; Araştırma Ormanı, 750 m., 19.07.2002 17♂8♀; Finike, Dağbağ, 475 m., 13.07.2002 1♀; Çığlık, 330 m., 19.07.2002 1♂; Korkuteli, Söğütcük, 1010 m., 13.07.2002 1♂1♀; Kaş, Kemerköy, 260 m., 13.07.2002 2♂1♀. It has been collected from weeds in *Pinus brutia* forest, on *Asphodelus aestivus*, *Verbascum*, *Phlomis*, *Fontanesia phillyreoides*, *Cytisus laburnum*, *Vitex agnus-castus* and *Cistus creticus*. **Distribution in Turkey:** Adana, Adiyaman, Afyon, Ankara, Aydin, Balikesir, Bilecik, Burdur, Bursa, Çanakkale, Çorum, Denizli, Eskisehir, Gaziantep, Gümüşhane, Hatay, İzmir, Kırklareli, Kırşehir, Kocaeli, Kütahya, Malatya, Manisa, Muğla, Sakarya, Sivas, Tekirdağ, Tokat, Uşak, Yozgat (Linnavuori, 1965; Dlabola, 1981; Lodos & Kalkandelen, 1981a; Kartal, 1985). **Remarks:** New for Antalya.

***Bubastia (Acrestia) suturalis* (Fieber, 1877)**

Materials: Manavgat, Bucakşeyhler, 45 m., 19.07.2001 3♂1♀; Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 1♂; Düzlerçami, 280 m., 22.07.2001 1♀; Güllük dağı, Araştırma Ormanı, 750 m., 19.07.2002 2♂; Çığlık, 330 m., 19.07.2002 3♂; Güllük Dağı, Araştırma Ormanı, 550 m., 19.07.2002 1♀; Kaş, Kemerköy, 260 m., 13.07.2002 1♀; Çamyuva, 20.08.2002 1♂1♀; Ovacık, 1150 m., 20.08.2002 1♀; İbradı, Akşahap-İbradı, 530 m., 21.08.2002 1♂; Manavgat, Beşkonak, Köprülü Kanyon, 495 m., 27.08.2003 1♂; Beşkonak, 180 m., 27.08.2003 1♂ on *Fontanesia phillyreoides*, *Quercus*, *Pistacia terebinthus* and *Myrthus communis*. **Distribution in Turkey:** Adana, Burdur, Hatay, İzmir, Muğla (Dlabola, 1957, Linnavuori, 1965; Lodos & Kalkandelen, 1981a). **Remarks:** New for Antalya.

Family: Flatidae Spinola, 1839

***Phantia subquadrata* (Herrich-Schäffer, 1838)**

Materials: Akseki, Güçlüköy, 610 m., 18.08.1999 1♀; Korkuteli, Kargalık, 1250 m., 27.07.1999 1♀; Akseki, Murtiçi, 540 m., 25.07.2001 12♂19♀; Termessos, Güllük Dağı, 540 m., 22.07.2001 1♀; Güllük Dağı, 900-1000 m., 22.07.2001 1♂; Araştırma Ormanı, 550 m., 19.07.2002 1♀; Araştırma Ormanı, 750 m.,

19.07.2002 1♂1♀; İbradı, Akşahap-İbradı, 530 m., 21.08.2002 1♀; Kemer, Sarıçınar Dağı, 1245 m., 20.08.2002 1♀. *Cytisus laburnum*, *Genista*, *Spartium junceum* and *Sesamum indicum*. **Distribution in Turkey:** Adana, Ankara, Aydin, Bilecik, Denizli, Diyarbakır, Edirne, Gaziantep, Isparta, İzmir, Konya, Manisa, Mardin, Muğla (Fahringer, 1922; Dlabola, 1957; Lodos & Kalkandelen, 1981b). **Remarks:** New for Antalya.

Family: Cicadidae Leach, 1815

***Lyristes plebejus* (Scopoli, 1763)**

Materials: Manavgat, Bucakşeyhler, 15-26.08.1997 3♂3♀; Demirciler, 65 m., 21.07.1999 1♂1♀; Akseki, Güneykaya, 570 m., 25.07.1999 1♂; Manavgat, Demirciler, 70 m., 22.07.1999 3♂; Akseki, Erenkaya, 680 m., 25.07.1999 3♂; Alanya, 24.07.1999 2♂; Korkuteli, Söğütçük, 570 m., 27.07.1999 1♂; Kaş, Beldibi, 1400 m., 29.07.1999 2♂1♀; Manavgat, Değirmenli, 26.07.1999 1♂; Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 1♀; Araştırma Ormanı, 550 m., 19.07.2002 1♂; Gündoğmuş, Güneycik, 210 m., 14.07.2002 4♂2♀; Kemer, Sarıçınar Dağı, 1245 m., 20.08.2002 1♂. On *Pinus brutia*, *Olea europaea*, *Amygdalus*, *Vitex agnus-castus*, *Quercus*, *Pistacia terebinthus*, *Arbustus* and maquis shrubs. **Distribution in Turkey:** Adana, Ankara, Antalya, Artvin, Aydin, Balikesir, Bursa, Denizli, Erzincan, Gaziantep, İzmir, İstanbul, Kırklareli, Manisa, Mardin, Muğla, Kütahya (Fahringer, 1922; Dlabola, 1957; Lodos & Kalkandelen, 1980b). **Remarks:** Known to Antalya.

***Cicadatra adanai* Kartal, 1980**

Materials: Gündoğmuş, Çiçekoluk, 960 m., 25.07.1999 1♀; Manavgat, Bucakşeyhler, 15-26.08.1997 1♂; Manavgat, 26.07.1999 3♂; İbradı, Üzümdere, 470 m., 18.08.1999 1♀; Manavgat, 30.07.1999 1♂; Demirciler, 40 m., 21.07.1999 1♂3♀; Demirciler, 70 m., 22.07.1999 1♂; Bucakşeyhler, 13.08.1999 1♂; Manavgat, 01.08.1999 1♂; Bucakşeyhler, 45 m., 19.07.2001 2♂1♀; Manavgat, 10 m., 26.07.2001 1♂; Gündoğmuş, Güneycik, 210 m., 22.06.2002 1♀; Güneycik, 210 m., 14.07.2002 1♂; Manavgat, Demirciler, 105 m., 21.06.2002 1♂; Güllük Dağı, Araştırma Ormanı, 550 m., 19.07.2002 1♀. On *Zea mays*, *Ficus carica*, *Olea europaea*, *Sesamum indicum*, *Fontanesia phillyreoides*, *Cistus creticus*, *Pinus brutia*, *Salix* and weeds. **Distribution in Turkey:** Adana (Kartal, 1980). **Remarks:** New for Antalya and endemic to Turkey.

***Cicadatra hyalina* (Fabricius, 1798)**

Materials: Manavgat, Demirciler, 40 m., 17.07.2002 1♀. On *Olea europaea*. **Distribution in Turkey:** Adiyaman, Ankara, Elazığ, Erzincan, Gümüşhane, Hakkari, Isparta, İzmir, Konya, Kahramanmaraş, Nevşehir, Niğde, Siirt, Urfa, Van (Dlabola, 1957; Lodos & Kalkandelen, 1981b; Kartal & Zeybekoğlu, 1999). **Remarks:** New for Antalya.

***Cicada mordoganensis* Boulard, 1979**

Materials: Manavgat, 29.07.1999 1♀; Demirciler, 70 m., 22.07.1999 13♂9♀; Demirciler, 65 m., 21.07.1999 7♂2♀; Değirmenli, 26.07.1999 1♂1♀; Alanya, Keşefli, 21.07.2001 7♂2♀; Manavgat, Demirciler, 105 m., 21.06.2002 2♂1♀. On *Olea europaea*, *Acacia*, and *Pinus brutia*. **Distribution in Turkey:** Antalya, İzmir, Muğla (Boulard, 1979; Lodos & Kalkandelen, 1981b). **Remarks:** It is only known from Antalya.

***Cicada lodosi* Boulard, 1979**

Materials: Düzlerçamı, 280 m., 22.07.2001 4♂1♀; Çığlık, 330 m., 19.07.2002 4♂3♀. *Pinus brutia* and *Olea europaea*. **Distribution in Turkey:** İzmir, Manisa (Boulard, 1979; Lodos & Kalkandelen, 1981b). **Remarks:** New for Antalya and endemic to Turkey.

***Cicada permagna* (Haupt, 1917)**

Materials: Manavgat, Bucakşeyhler, 18-28.06.1997 6♂3♀; Bucakşeyhler, 45 m., 19.07.2001 5♂; Akseki, Erenkaya, 630 m., 22.06.2002 1♂; Güllük Dağı, Araştırma Ormanı, 750 m., 19.07.2002 1♂1♀; Gündoğmuş, Çiçekoluk, 960 m., 14.07.2002 1♀; Kemer, Olimpos, 23.07.2003 1♂. On *Pinus brutia* and *Olea europaea*. **Distribution in Turkey:** Antalya, Mersin (Boulard, 1979; Dlabola, 1981). **Remarks:** Endemic to Turkey and known to Antalya.

***Tibicina haematodes* (Scopoli, 1763)**

Materials: Kaş, Beldibi, 1400 m., 29.07.1999 1♂. On *Pinus brutia*. **Distribution in Turkey:** İstanbul (Fahringer, 1922). **Remarks:** New for Antalya.

***Pagiphora aschei* Kartal, 1978**

Materials: Manavgat, Demirciler, 40 m., 21.07.1999 1♀; Akseki, Güneykaya, 570 m., 25.07.1999 1♂; Termessos, Güllük Dağı, 900-1000 m., 22.07.2001 1♀; Güllük dağı, 540 m., 22.07.2001 1♂1♀; Araştırma Ormanı, 550 m., 19.07.2002 1♀; Araştırma Ormanı, 750 m., 19.07.2002 1♂1♀; Manavgat, Demirciler, 17.07.2002 1♂. On *Ficus carica*, *Quercus*, *Arbutus andrachne* and *Ulmus*. **Distribution in Turkey:** Antalya (Kartal, 1983). **Remarks:** Known to Antalya.

***Cicadetta dimissa* (Hagen, 1856)**

Materials: Manavgat, Bucakşeyhler, 18-28.06.1997 1♂. On shrubs. **Distribution in Turkey:** Antalya, Aydın, Gümüşhane, Isparta, İzmir (Lodos & Kalkandelen, 1981b). **Remarks:** Known to Antalya.

Family: Cercopidae Leach, 1815***Triecphorella geniculata* (Horvath, 1881)**

Materials: Gündoğmuş, Güneycik, 210 m., 17.03.2002 1♀; Güneycik, 210 m., 21.04.2002 1♀; Serinyaka, 640 m., 19.05.2002 1♀. It has been collected from weeds in *Pinus brutia* forest and *Myrthus communis*. **Distribution in Turkey:** Adana, Kahramanmaraş, Konya, Samsun (Fahringer, 1922; Kartal, et al., 1994). **Remarks:** New for Antalya.

***Lepyronia coleoptrata* (Linnaeus, 1758)**

Materials: Manavgat, Değirmenli, 04.08.1999 2♂2♀; Bucakşeyhler, 13.08.1999 11♂7♀; Demirciler, 10 m., 17.08.1999 2♂2♀ Bucakşeyhler, 24.07.1999 5♂2♀; Bucakşeyhler, 30 m., 23.07.1999 1♂1♀; Manavgat, 10.08.1999 1♂; Demirciler, 65 m., 21.07.1999 1♀; Bucakşeyhler, 45 m., 24.07.1999 1♂2♀; Aksu, Perge, 27.07.1999 1♂1♀; Manavgat, 10 m., 20.07.1999 1♀; Demirciler, Külcüler, 150 m., 22.07.1999 1♂; Alanya, Okurcalar, 12.05.2001 1♂; Manavgat, Tilkiler, 220 m., 14.05.2001 2♂2♀; Demirciler, 40 m., 11.05.2001 1♂3♀; Demirciler, 40 m., 05.06.2001 1♂6♀; Finike, 06.06.2001 1♂; Gündoğmuş, Güneycik, 210 m., 08.06.2001 1♀; Manavgat, Bucakşeyhler, 45 m., 19.07.2001 1♂; Demirciler, 40 m., 24.07.2001 1♀; Alanya, Keşfeli, 21.07.2001

1♀; Yeşilköy, 20.05.2002 2♂3♀; Okurcalar, 20.05.2002 3♀; Gazipaşa, Çalıpinar-Kahyalar, 20 m., 20.05.2002 1♂1♀; Manavgat, Demirciler, 20 m., 18.05.2002 2♂; Demirciler 40 m., 20.06.2002 3♂6♀; Korkuteli, Sülekler, 1300 m., 23.06.2002 3♂1♀; Koru Dağ, Kartal Yaylası, 1600 m., 23.06.2002 4♂3♀; Manavgat, Demirciler, 65 m., 15.07.2002 1♂; Finike, 13.07.2002 1♀; Gazipaşa, Macarköy, 22.08.2002 2♀. It has been collected from weeds in *Pinus brutia* forest, on *Populus*, *Salix*, *Verbascum* and maquis shrubs. **Distribution in Turkey:** Adana, Afyon, Ankara, Artvin, Aydin, Bilecik, Bursa, Çanakkale, Çankırı, Çorum, Diyarbakır, Edirne, Erzincan, Gümüşhane, İzmir, Kars, Kütahya, Manisa, Mardin, Muğla, Muş, Sakarya, Samsun, Siirt, Tokat (Fahringer, 1922; Dlabola, 1957; Lodos & Kalkandelen, 1981c; Kartal, et. al., 1994). **Remarks:** New for Antalya.

***Neophilaenus campestris* (Fallen, 1805)**

Materials: Manavgat-Çeltikçi, 14.04.2001 1♀; Belenobası, 130 m., 15.04.2001 1♂5♀; Alanya, Okurcalar, 12.05.2001 3♀; Keşefli, 12.05.2001 8♂18♀; Gazipaşa, Çalıpinar, 25 m., 12.05.2001 2♂; Manavgat, Yukarıışıklar-Tilkiler, 460 m., 14.05.2001 1♂3♀; Tilkiler, 220 m., 14.05.2001 2♂1♀; Yaylaalan, 540 m., 14.05.2001 1♀; Tilkiler, 470 m., 14.05.2001 1♂1♀; Demirciler, 40 m., 11.05.2001 1♂; Bucakşeyhler, Seleukeia, 250 m., 23.04.2002 2♀; Demirciler, 105 m., 22.04.2002 1♂2♀; Demirciler, 40 m., 22.04.2002 1♀; Gündoğmuş, Güneycik, 240 m., 21.04.2002 1♂; Çiçekoluk, 960 m., 19.05.2002 3♂1♀; Alanya, Okurcalar, 20.05.2002 2♀; Gazipaşa, Çalıpinar-Kahyalar, 20 m., 20.05.2002 1♂3♀. It has been collected from weeds in *Pinus brutia* forest and fields. **Distribution in Turkey:** Adana, Adiyaman, Ankara, Artvin, Bitlis, Bursa, Çanakkale, Eskişehir, Giresun, Hatay, İzmir, Kırklareli, Kütahya, Manisa, Muğla, Nevşehir, Samsun, Siirt, Sinop, Trabzon, Van (Linnauvori, 1965; Lodos & Kalkandelen, 1981c). **Remarks:** New for Antalya.

***Neophilaenus minor* (Kirschbaum, 1868)**

Materials: Korkuteli, Koru Dağı, Kartal Yaylası, 1600 m., 23.06.2002 4♂4♀. It has been collected from weeds in *Juniperus* forest. **Distribution in Turkey:** Ankara, Konya, Sivas (Dlabola, 1957, 1981). **Remarks:** New for Antalya.

***Aphrophora corticea* Germar, 1821**

Materials: Manavgat, Yaylaalan, 545 m., 14.05.2001 1♂; Yukarıışıklar-Tilkiler, 460 m., 14.05.2001 1♂3♀; Gündoğmuş, Serinyaka, 640 m., 19.05.2002 3♀; Güneycik, 240 m., 19.05.2002 1♀; Akseki, Güçlüköy, 575 m., 22.06.2002 1♀. It has been collected from weeds and shrubs in *Pinus brutia* forest and on *P. brutia*. **Distribution in Turkey:** Afyon, Aydin, Bilecik, Çanakkale, Isparta, Kütahya, Muğla, Tokat. (Lodos & Kalkandelen, 1981c). **Remarks:** New for Antalya.

***Philaenus signatus* Melchar, 1896**

Materials: Termessos, Güllük Dağı, 900-1000 m., 21.05.2002 1♂. On shrubs. **Distribution in Turkey:** Adana, İzmir, Kahramanmaraş, Muğla (Dlabola, 1957, 1971; Lodos & Kalkandelen, 1981c). **Remarks:** New for Antalya.

***Philaenus spumarius* (Linnaeus, 1758)**

Materials: Gündoğmuş, Güneycik, 210 m., 17.04.2001 2♂3♀; Manavgat, Demirciler, Külcüler, 170 m., 15.04.2001 7♂8♀; Manavgat-Çeltikçi, 14.04.2001 1♂; Belenobası, 130 m., 15.04.2001 13♂17♀; Demirciler, 65 m., 14.04.2001

6♂8♀; Saraklı, 15.04.2001 1♂1♀; Hocalı, 215 m., 15.04.2001 2♂; Demirciler, 40 m., 14..04.2001 3♀; Yaylaalan, 540 m., 14.05.2001 25♂17♀; Tilkiler, 470 m., 14.05.2001 7♂2♀; Demirciler, 65 m., 11.05.2001 14♂25♀; Alanya, Okurcalar, 12.05.2001 4♂5♀; Keşfili, 12.05.2001 6♀; Manavgat, Tilkiler, 220 m., 14.05.2001 21♂18♀; Yaylaalan, 545 m., 14.05.2001 3♂3♀; Gazipaşa, Çalıpınar, 25 m., 12.05.2001 3♂2♀; Yukarışıkalar-Tilkiler, 460 m., 14.05.2001 38♂51♀; Beşkonak, 145 m., 07.06.2001 1♂; Demirciler, 40 m., 05.06.2001 1♀; Evrenleryavşı, Söylerkale, 250 m., 16.03.2002 1♀; Demirciler, 40 m., 22.04.2002 5♂7♀; Bucakşeyhler, 70 m., 23.04.2002 9♂6♀; Bucakşeyhler, Seleukeia, 250 m., 23.04.2002 2♂3♀; Demirciler, Kılçüler, 230 m., 22.04.2002 2♂1♀; Demirciler, 105 m., 22.04.2002 2♂3♀; Demirciler, 65 m., 22.04.2002 14♂12♀; Gündoğmuş, Güneycik, 240 m., 21.04.2002 2♂1♀; Serinyaka, 640 m., 19.05.2002 15♂5♀; Çiçekoluk, 960 m., 19.05.2002 3♂; Manavgat, Demirciler, 20 m., 18.05.2002 1♂3♀; Demirciler, 105 m., 18.05.2002 3♂; Gazipaşa, Çalıpınar-Kahyalar, 20.05.2002 2♂3♀; Alanya, Okurcalar, 20.05.2002 1♂; Akseki, Güçlüköy, 575 m., 22.06.2002 1♀. It has been collected from weeds in *Pinus brutia* forest, on *Myrthus communis*, *Pistacia terebinthus*, *Phlomis* and fields. **Distribution in Turkey:** Ağrı, Amasya, Ankara, Artvin, Aydın, Balıkesir, Bilecik, Bitlis, Bursa, Çanakkale, Elazığ, Erzincan, Erzurum, Giresun, Gümüşhane, Hakkari, İzmir, Kahramanmaraş, Kars, Kocaeli, Kırklareli, Kütahya, Malatya, Manisa, Mardin, Muğla, Ordu, Rize, Samsun, Siirt, Sinop, Sivas, Tekirdağ, Tokat, Trabzon, Van (Dlabola, 1957, 1981; Linnauvori, 1965; Lodos & Kalkandelen, 1981c; Kartal, et. al., 1994). **Remarks:** New for Antalya.

Family: Membracidae Rafinesque, 1815

***Gargara genistae* (Fabricius, 1775)**

Materials: Termessos, Güllük Dağı, 540 m., 22.07.2001 5♀; Güllük Dağı, 900-1000 m., 22.07.2001 2♂8♀; Güllük Dağı, 900-1000 m., 19.07.2002 7♂5♀; Araştırma Ormanı, 550 m., 19.07.2002 8♀; Araştırma Ormanı, 750 m., 19.07.2002 19♀; Çığlık, 330 m., 19.07.2002 3♀. On Fabaceae shrubs and *Cytisus laburnum*. **Distribution in Turkey:** Ankara, Aydın, Balıkesir, Çorum, Erzurum, İzmir, Muğla (Dlabola, 1957; Lodos & Kalkandelen, 1981c). **Remarks:** New for Antalya.

***Centrotus cornutus* (Linnaeus, 1758)**

Materials: Termessos, Güllük Dağı, 900-1000 m., 21.05.2002 1♂1♀. On Fabaceae shrubs. **Distribution in Turkey:** Artvin, Aydın, Bilecik, Bursa, Çorum, Denizli, Erzincan, Edirne, Giresun, İstanbul, İzmir, Kocaeli, Kütahya, Kırklareli, Ordu, Sakarya, Tekirdağ (Fahringer, 1922; Lodos & Kalkandelen, 1981c). **Remarks:** New for Antalya.

* This study is prepared from a Ph. D. Thesis approved by the Institute of Natural Sciences of Gazi University on August 2005.

ACKNOWLEDGEMENTS

I thank Dr. Yasemin Özdemir, Dr. Mustafa Özdemir, Dr. İşıl Özdemir and my family for their help during different phases of this project.

LITERATURE CITED

- Akman, Y.** 1995. Türkiye Orman Vejetasyonu. Ankara Üniversitesi Fen Fakültesi Yayınları, Ankara 326 pp.
- Asche, M.** 1982. Beiträge zur Delphaciden-Fauna der Türkei (Anatolien) (Homoptera Cicadina Delphacidae). Marburger Entomologische Publikationen 1(7): 71-98.
- Boulard, M.** 1979. Cigales du genre *Cicada* Linne, originaires de Turquie (Hom. Cicadidae). Türkiye Bitki Koruma Dergisi 3(2): 67-74.
- Boulard, M.** 1993. *Pagiphora yanni*, Cigale anatoliennes inédites. Description et premières informations biologiques (cartes d'identité et d'éthologie sonores) (Homoptera, Cicadoidea, Tibicinidae). Türkiye Entomoloji Dergisi 17(1): 1-9.
- Demir, E.** 1998. Türkiye Homoptera Faunası İçin Yeni Kayıtlar (Homoptera, Auchenorrhyncha, Tropiduchidae). Centre Entomological Studies Ankara, Miscellaneous Papers 55/56: 1-3.
- Demir, E.** 2006. Contributions to the knowledge of Turkish Auchenorrhyncha (Homoptera) with a new record, *Pentastiridius nanus* (Ivanov, 1885). Munis entomology and zoology 1(1): 97-122.
- Dlabola, J.** 1957. Results of the Zoological Expedition of the National Museum in Prague to Turkey 20. Homoptera, Auchenorrhyncha. Acta Entomologica Musei Nationalis Pragae 31(469): 19-68.
- Dlabola, J.** 1971a. Taxonomische und Chorologische Ergänzungen der Zikadenfauna von Anatolien, Iran, Afghanistan und Pakistan (Homoptera, Auchenorrhyncha). Acta entomologica Bohemoslovaca 68: 377-396.
- Dlabola, J.** 1971b. Taxonomische und chorologische Erzangungen zur türkischen und iranischen Zikadenfauna (Homoptera, Auchenorrhyncha). Acta faunistica entomologica Musei Nationalis Pragae 14(163): 115-138.
- Dlabola, J.** 1974. Zur Taxonomie und Chorologie einiger Mediterraner Zikaden (Homoptera, Auchenorrhyncha). Acta Zoolojica Academiae Scientiarum Hungaricae 20(3-4): 289-308.
- Dlabola, J.** 1977. Neue Zikaden-Taxone von *Mycterodus*, *Erythria*, *Selenocephalus* und *Goldeus* (Homoptera, Auchenorrhyncha). Acta Zoolojica Academiae Scientiarum Hungaricae 23(3-4): 279-292.
- Dlabola, J.** 1979a. Neue Zikaden aus Anatolien, Iran und aus Südeuropäischen Landern (Homoptera, Auchenorrhyncha). Acta Zoolojica Academiae Scientiarum Hungaricae 25(3-4): 235-257.
- Dlabola, J.** 1979b. *Tshurtshurnella*, *Bubastia* und andere verwandte Taxone (Auchenorrhyncha, Issidae). Acta entomologica Bohemoslovaca 76: 266-286.
- Dlabola, J.** 1980. Tribus-Einteilung, neue Gattungen und Arten der subf. Issinae in der Eremischen Zone (Homoptera Auchenorrhyncha). Sbornik Narodniho Muzea V Praze 36(4): 173-248.
- Dlabola, J.** 1981. Ergebnisse der Tschechoslowakisch-Iranischen Entomologischen Expeditionen nach dem Iran (1970 und 1973). Acta Entomologica Musei Nationalis Pragae 40: 127-311.

- Dlabola, J.** 1982. Fortsetzung der Ergänzungen zur Issiden-Taxonomie von Anatolien, Iran und Griechenland (Homoptera, Auchenorrhyncha). Sbornik Narodniho Muzea V Praze 38(3): 113-169.
- Dlabola, J.** 1983a. Ergebnisse der Tschechoslovakisch-Iranischen Entomologischen Expeditionen 1970 und 1973 nach dem Iran. Acta Entomologica Musei Nationalis Pragae 41: 91-97.
- Dlabola, J.** 1983b. Neue mediterrane, meistens anatolische Issidaen (Homoptera, Auchenorrhyncha). Acta entomologica Bohemoslovaca 80:114-136.
- Dlabola, J.** 1984a. Neue Mediterrane, Eremische und Ostafrikanische Issiden-Taxone (Homoptera, Auchenorrhyncha). Sbornik Narodniho Muzea V Praze 40(3-4): 217-243.
- Dlabola, J.** 1984b. Typenrevision einiger mediterraner bzw. Nordafrikanischer Hysteropterum (s.l.) Auchenorrhyncha-Issidae. Acta faunistica entomologica Musei Nationalis Pragae 17(195): 27-68.
- Dlabola, J.** 1985. Neue Cixiide vom Iran, Nachbarlandern und anderen Mediterrangebieten (Homoptera, Auchenorrhyncha). Acta entomologica Bohemoslovaca 82(2): 95-128.
- Dlabola, J.** 1986. Neue Arten der Fulgoromorphen Zikaden-Familien vom Mittelmeergebiet und nahen Osten. Sbornik Narodniho Muzea V Praze 42(3-4): 169-196.
- Dlabola, J.** 1995. Mycterus verwandte Taxone und sieben neue Zikadenarten (Homoptera, Auchenorrhyncha). Acta Entomologica Musei Nationalis Pragae 44: 301-319.
- Drosopoulos, S.** 2003. New data on the nature and origin of colour polymorphism in the spittlebug genus *Philaenus* (Hemiptera, Aphrophoridae), Annales de la Société entomologique de France (NS) 39: 31-42.
- Drosopoulos, S., Asche, M. & Hoch, H.** 1985. Contribution to the planthopper fauna of Greece (Homoptera, Auchenorrhyncha, Fulgoromorpha, Delphacidae), Annals of the Benaki Phytopathological Institute: 35-88.
- Drosopoulos, S., Eliopoulos, E. & Tsakalou, P.** 2006. Is Migration Responsible for the Peculiar Geographical Distribution and Speciation Based on Acoustic Divergence of Two Cicadas in the Aegean Archipelago? In *Insect Sounds and Communication. Physiology, Behaviour, Ecology and Evolution*, Drosopoulos, S and Claridge, M. F. , Eds., CRC Press, Taylor and Francis, London, pp. 219-226
- D'Urso, V. & Guglielmino, A.** 1995. Taxonomic remarks on Italian *Cixidia* with description of two new species (Insecta, Homoptera, Auchenorrhyncha, Achilidae). Spixiana, 18 (1): 49-64.
- Emeljanov, A.F.** 2002. New taxa and new data on distribution of the subfamily Orgeriinae in the Mediterranean (Homoptera: Dictyopharidae). Zoosystematica Rossica 11(2): 311-319.
- Emeljanov, A. F.** 2004. The subgeneric division of the genus *Dictyophara* Germar, 1833 (Homoptera: Dictyopharidae). Russian Entomological Journal, 12 (4): 357-358.
- Fahringer, J.** 1922. Eine Rhynchotenausbeute aus der Türkei, Kleinasien und den benachbarten Gebieten. Konowia 1: 296-307.

- Gnezdilov, V.M.** 2002. New species of the genus *Tshurtshurnella* Kusnezov, 1927 (Homoptera: Cicadina, Issidae) from Turkey and Lebanon. Russian Entomological Journal 11(3): 233-240.
- Gnezdilov, V. M. & Mazzoni, V.** 2004. Notes on the *Latilica maculipes* (Melichar, 1906) species group (Homoptera, Issidae). Redia, 86, 2003:147-151.
- Gnezdilov, V. M., Drosopoulos, S. & Wilson, M. R.** 2004. New data on the taxonomy and distribution of soma Fulgoroidea (Homoptera, Cicadina). Zoosystematica Rossica, 12 (2), 2003: 217-223.
- Güçlü, Ş.** 1996. Studies on Delphacidae (Hom., Auchenorrhyncha) from Turkey. Turkish Journal of Zoology 20: 407-411.
- Hoch, H. & Remane, R.** 1985. Evolution und Speziation der Zikaden-Gattung *Hyalesthes* Signoret, 1865 (Homoptera, Auchenorrhyncha, Fulgoroidea, Cixiidae). Marburger Entomologische Publikationen 2(2): 1-427.
- Huang, J. & Bourgoin, T.** 1993 The planthopper genus *Trypetimorpha*: systematics and phylogenetics relationships (Hemiptera: Fulgoromorpha: Tropiduchiae). J. Nat. Hist., 27: 609-629.
- Kalkandelen, A.** 1980. Contributions to the families Delphacidae and Cicadellidae (Homoptera) from Turkey. Türkiye Bitki Koruma Dergisi 4(3): 147-154.
- Kalkandelen, A.** 1987-2000. Türkiye Cixiidae (Homoptera) türleri üzerine taksonomik çalışmalar. Bitki Koruma Bülteni 27(3-4): 119-146; 28(3-4), 113-140; 29(1-4), 1-17, 117-132; 30(1-4), 3-27; 33(3-4), 65-82; 34(1-2), 1-21; 40(3-4). 91-123.
- Kartal, V.** 1978. Eine neue Singzikaden Art-der Gattung *Pagiphora* Horvath von Kreta (Homoptera, Auchenorrhyncha, Cicadidae). Entomologische Zeitschrift 88(16): 179-185.
- Kartal, V.** 1980. Neue und wenig bekannte Arten der Gattungen Cicadetta und Cicadatra aus dem Irak und der Türkei (Homoptera, Auchenorrhyncha, Cicadidae). Entomologische Zeitschrift 17: 185-192.
- Kartal, V.** 1981. Neue Homopteren aus der Türkei-I. Priamus 1(1): 24-30.
- Kartal, V.** 1983. Neue Homopteren aus der Türkei II (Homoptera Auchenorrhyncha). Marburger Entomologische Publikationen 1(8): 235-248.
- Kartal, V.** 1985a. Neue und wenig bekannte *Tshurtshurnella*-Arten (Homoptera Auchenorrhyncha Issidae) aus der Türkei. Marburger Entomologische Publikationen 1(10): 191-218.
- Kartal, V.** 1985b. Türkiye Yukarı Kızılırmak Havzasındaki Issidae (Homoptera, Auchenorrhyncha) Familyası Türlerinin Taksonomik Yönden İncelenmesi. Doğa A2: 64-77.
- Kartal, V.** 1986. Wenig bekannte und für die Türkei neue *Nymphorgerius*-Arten (Homoptera, Auchenorrhyncha, Dictyopharidae). Türkiye Bitki Koruma Dergisi 10(2): 99-103.
- Kartal, V.** 1987. Eine neue und wenig bekannte Arten der Gattung *Ranissus* aus der Türkei (Homoptera, Auchenorrhyncha, Dictyopharidae). Türkiye Entomoloji Dergisi 11(3): 145-150.

Kartal, V., Zeybekoğlu, Ü. & Özdemir, G. 1994. Samsun Çevresinde Cercopidae (Hom., Auchenorrhyncha) Familyası Türleri Üzerine Taksonomik Bir Araştırma. Ondokuz Mayıs Üniversitesi Fen Dergisi 5(1): 147-157.

Linnavuori, R. 1965. Studies on the South and East-Mediterranean Hemipterous fauna. Acta Entomologica Fennica, Helsinki 21: 1-70.

Lodos, N. & Kalkandelen, A. 1980a. Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey I. Family Cixiidae Spinola. Türkiye Bitki Koruma Dergisi 4(1): 15-27.

Lodos, N. & Kalkandelen, A. 1980b. Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey II. Family Delpacidae Leach. Türkiye Bitki Koruma Dergisi 4(2): 103-117.

Lodos, N. & Kalkandelen, A. 1980c. Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey III. Families Meenoplidae, Derbidae, Achilidae, Dictyopharidae and Tettigometridae. Türkiye Bitki Koruma Dergisi 4(3): 161-178.

Lodos, N. & Kalkandelen, A. 1981a. Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey IV. Family Issidae Spinola. Türkiye Bitki Koruma Dergisi 5(1): 5-21.

Lodos, N. & Kalkandelen, A. 1981b. Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey V. Families Flatidae, Ricanidae and Cicadidae. Türkiye Bitki Koruma Dergisi 5(2): 67-82.

Lodos, N. & Kalkandelen, A. 1981c. Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey VI. Families Cercopidae and Membracidae. Türkiye Bitki Koruma Dergisi 5(3): 133-149.

Lodos, N. & Kalkandelen, A. 1988. Preliminary list of Auchenorrhyncha with notes on distribution and importance of Turkey XXVII. (Addenda and Corrigenda). Türkiye Entomoloji Dergisi 12(1): 11-22.

Quartau, J. A., Rebelo, M. T., Simões, P. C., Fernandes, T. M., Claridge, M. F., Drosopoulos, S. & Morgan, J. C. 1999. Acoustic signals of populations of *Cicada orni* L. in Portugal and Greece (Hemiptera: Auchenorrhyncha: Cicadomorpha: Cicadidae). Reichenbachia Staatliches Museum für Tierkunde Dresden. 33(8): 71-80.

Quartau, J. A. & Simões, P. C. 2006. Acoustic Evolutionary Divergence in Cicadas: The Species of *Cicada* L. in Southern Europe. In *Insect Sounds and Communication. Physiology, Behaviour, Ecology and Evolution*, Drosopoulos, S and Claridge, M. F. , Eds., CRC Press, Taylor and Francis, London, pp. 227-237